


Module 3: Duality through examples

Recap: Shortest Paths

In an instance of the **shortest path** problem, we are given

- a **graph** $G = (V, E)$, a non-negative length c_e for each edge $e \in E$, and
- a pair of vertices s and t in V .




Recap: Shortest Paths

In an instance of the **shortest path** problem, we are given

- a **graph** $G = (V, E)$, a non-negative length c_e for each edge $e \in E$, and
- a pair of vertices s and t in V .


Our **goal** is to compute an s, t -**path** P of smallest total length.



Recap: Shortest Paths

In an instance of the **shortest path** problem, we are given

- a **graph** $G = (V, E)$, a non-negative length c_e for each edge $e \in E$, and
- a pair of vertices s and t in V .



Our **goal** is to compute an **s, t -path** P of smallest total length.

Recall: an s, t -path is a sequence

$$P := u_1 u_2, u_2 u_3, \dots, u_{k-1} u_k$$

where

- $u_i u_{i+1} \in E$ for all i , and
- $u_1 = s$, $u_k = t$, and $u_i \neq u_j$ for all $i \neq j$.

Recap: Shortest Paths

In an instance of the **shortest path** problem, we are given

- a **graph** $G = (V, E)$, a non-negative length c_e for each edge $e \in E$, and
- a pair of vertices s and t in V .


Our **goal** is to compute an s, t -path P of smallest total length.

Recall: an s, t -path is a sequence

$$P := u_1 u_2, u_2 u_3, \dots, u_{k-1} u_k$$

where

- $u_i u_{i+1} \in E$ for all i , and
- $u_1 = s$, $u_k = t$, and $u_i \neq u_j$ for all $i \neq j$.




Its **length** is given by

$$c(P) = c_{u_1 u_2} + c_{u_2 u_3} + \dots + c_{u_{k-1} u_k}$$

In the example, we see by inspection that

$$P = sa, ac, cb, bt$$


is a **shortest path** and that its length is 9.



In the example, we see by inspection that

$$P = sa, ac, cb, bt$$

is a **shortest path** and that its length is 9.




Question

1. Given a shortest-path instance and a candidate shortest s, t -path P , is there a **short proof** of its optimality?

In the example, we see by inspection that

$$P = sa, ac, cb, bt$$

is a **shortest path** and that its length is 9.




Question

1. Given a shortest-path instance and a candidate shortest s, t -path P , is there a **short proof** of its optimality?
2. How can we **find** a shortest s, t -path?

In the example, we see by inspection that

$$P = sa, ac, cb, bt$$

is a **shortest path** and that its length is 9.



Question


1. Given a shortest-path instance and a candidate shortest s, t -path P , is there a **short proof of its optimality**?
2. How can we **find** a shortest s, t -path?

We will answer both questions in this module. This lecture focus on question 1.

Shortest Paths: Finding an Intuitive Lower Bound

Cardinality Case

To make our lives easier, we will first consider the **cardinality special case** of the shortest path problem.




Cardinality Case

To make our lives easier, we will first consider the **cardinality special case** of the shortest path problem.

We consider shortest path instances where...

- each edge $e \in E$ has **length 1**,
and




Cardinality Case

To make our lives easier, we will first consider the **cardinality special case** of the shortest path problem.

We consider shortest path instances where...

- each edge $e \in E$ has **length 1**, and
- we are therefore looking for an s, t -path with the **smallest number of edges**.




Cardinality Case

To make our lives easier, we will first consider the **cardinality special case** of the shortest path problem.

We consider shortest path instances where...

- each edge $e \in E$ has **length 1**, and
- we are therefore looking for an s, t -path with the **smallest number of edges**.



Example: In the diagram above, one easily sees that

$$P = sj, ji, ig, gt$$

is a **shortest s, t -path**.


How can we prove this fact?

Cardinality Case

To make our lives easier, we will first consider the **cardinality special case** of the shortest path problem.

We consider shortest path instances where...

- each edge $e \in E$ has **length 1**, and
- we are therefore looking for an s, t -path with the **smallest number of edges**.



Example: In the diagram above, one easily sees that

$$P = sj, ji, ig, gt$$

is a **shortest s, t -path**.

How can we prove this fact?

→ The answer lies in **s, t -cuts!**

Definition

For $U \subseteq V$, we define

$$\delta(U) = \{uv \in E : u \in U, v \notin U\}$$

and call it an s, t -cut if $s \in U$, and $t \notin U$.

s, t -cuts

Definition

For $U \subseteq V$, we define

$$\delta(U) = \{uv \in E : u \in U, v \notin U\}$$


and call it an s, t -cut if $s \in U$, and $t \notin U$.

Example

Let $U = \{s, a, j\}$. It follows that

$$\delta(U) = \{ab, ah, ji\}$$

is an s, t -cut.



Definition

For $U \subseteq V$, we define

$$\delta(U) = \{uv \in E : u \in U, v \notin U\}$$

and call it an s, t -cut if $s \in U$, and $t \notin U$.

Recall:


- If P is an s, t -path and $\delta(U)$ an s, t -cut, then P contains an edge of $\delta(U)$.

Example

Let $U = \{s, a, j\}$. It follows that

$$\delta(U) = \{ab, ah, ji\}$$

is an s, t -cut.



Definition

For $U \subseteq V$, we define

$$\delta(U) = \{uv \in E : u \in U, v \notin U\}$$

and call it an s, t -cut if $s \in U$, and $t \notin U$.

Recall:


- If P is an s, t -path and $\delta(U)$ an s, t -cut, then P contains an edge of $\delta(U)$.
- If $S \subseteq E$ contains an edge from every s, t -cut, then S contains an s, t -path.

Example

Let $U = \{s, a, j\}$. It follows that


$$\delta(U) = \{ab, ah, ji\}$$

is an s, t -cut.



From Cuts to Lower-Bounds

The example on the right shows 4 s, t -cuts, $\delta(U_1), \delta(U_2), \delta(U_3), \delta(U_4)$.



$$\delta(U_1) = \{sa, sj\}$$

$$\delta(U_2) = \{ab, ah, ji\}$$

$$\delta(U_3) = \{bc, hc, ig\}$$


$$\delta(U_4) = \{dt, gt\}$$

From Cuts to Lower-Bounds

The example on the right shows 4 s, t -cuts, $\delta(U_1), \delta(U_2), \delta(U_3), \delta(U_4)$.

Two important notes:

(1) $\delta(U_i) \cap \delta(U_j) = \emptyset$ for $i \neq j$ and



$$\delta(U_1) = \{sa, sj\}$$

$$\delta(U_2) = \{ab, ah, ji\}$$

$$\delta(U_3) = \{bc, hc, ig\}$$


$$\delta(U_4) = \{dt, gt\}$$

From Cuts to Lower-Bounds

The example on the right shows 4 s, t -cuts, $\delta(U_1), \delta(U_2), \delta(U_3), \delta(U_4)$.

Two important notes:

- (1) $\delta(U_i) \cap \delta(U_j) = \emptyset$ for $i \neq j$ and
- (2) an s, t -path must contain an edge from $\delta(U_i)$ for all i .



$$\delta(U_1) = \{sa, sj\}$$

$$\delta(U_2) = \{ab, ah, ji\}$$

$$\delta(U_3) = \{bc, hc, ig\}$$


$$\delta(U_4) = \{dt, gt\}$$

From Cuts to Lower-Bounds

The example on the right shows 4 s, t -cuts, $\delta(U_1), \delta(U_2), \delta(U_3), \delta(U_4)$.

Two important notes:

- (1) $\delta(U_i) \cap \delta(U_j) = \emptyset$ for $i \neq j$ and
- (2) an s, t -path must contain an edge from $\delta(U_i)$ for all i .



→ Every s, t -path must have at least 4 edges.

$$\delta(U_1) = \{sa, sj\}$$

$$\delta(U_2) = \{ab, ah, ji\}$$

$$\delta(U_3) = \{bc, hc, ig\}$$

$$\delta(U_4) = \{dt, gt\}$$

From Cuts to Lower-Bounds


The example on the right shows 4 s, t -cuts, $\delta(U_1), \delta(U_2), \delta(U_3), \delta(U_4)$.

Two important notes:

- (1) $\delta(U_i) \cap \delta(U_j) = \emptyset$ for $i \neq j$ and
- (2) an s, t -path must contain an edge from $\delta(U_i)$ for all i .

→ Every s, t -path must have at least 4 edges.

→ sj, ji, ig, gt is a shortest s, t -path!



$$\delta(U_1) = \{sa, sj\}$$

$$\delta(U_2) = \{ab, ah, ji\}$$


$$\delta(U_3) = \{bc, hc, ig\}$$

$$\delta(U_4) = \{dt, gt\}$$

From Cuts to Lower-Bounds

Question

Notice: hi is not in any of the $\delta(U_i)$. Does this mean that hi is not on any **shortest** s, t -path?



$$\delta(U_1) = \{sa, sj\}$$

$$\delta(U_2) = \{ab, ah, ji\}$$


$$\delta(U_3) = \{bc, hc, ig\}$$

$$\delta(U_4) = \{dt, gt\}$$

From Cuts to Lower-Bounds

Question

Notice: hi is not in any of the $\delta(U_i)$. Does this mean that hi is not on any **shortest** s, t -path?



Yes!

An s, t -path that contains hi must also contain an edge from **each** of the s, t -cuts $\delta(U_i)$.

$$\delta(U_1) = \{sa, sj\}$$

$$\delta(U_2) = \{ab, ah, ji\}$$


$$\delta(U_3) = \{bc, hc, ig\}$$

$$\delta(U_4) = \{dt, gt\}$$

From Cuts to Lower-Bounds

Question

Notice: hi is not in any of the $\delta(U_i)$. Does this mean that hi is not on any **shortest** s, t -path?



Yes!

An s, t -path that contains hi must also contain an edge from **each** of the s, t -cuts $\delta(U_i)$. \rightarrow It must contain **at least 5 edges!**

$$\delta(U_1) = \{sa, sj\}$$


$$\delta(U_2) = \{ab, ah, ji\}$$

$$\delta(U_3) = \{bc, hc, ig\}$$

$$\delta(U_4) = \{dt, gt\}$$

Back to the General Case

In general instances, we assign a **non-negative width** y_U to every s, t -cut $\delta(U)$.




Back to the General Case

In general instances, we assign a **non-negative width** y_U to every s, t -cut $\delta(U)$.

Definition

A width assignment $\{y_U : \delta(U) \text{ } s, t\text{-cut}\}$ is **feasible** if,




Back to the General Case

In general instances, we assign a **non-negative width** y_U to every s, t -cut $\delta(U)$.

Definition

A width assignment $\{y_U : \delta(U) \text{ } s, t\text{-cut}\}$ is **feasible** if, for every edge $e \in E$,




Back to the General Case

In general instances, we assign a **non-negative width** y_U to every s, t -cut $\delta(U)$.

Definition

A width assignment $\{y_U : \delta(U) \text{ } s, t\text{-cut}\}$ is **feasible** if, for every edge $e \in E$, the **total width** of all cuts containing e is no more than c_e .




Back to the General Case

In general instances, we assign a **non-negative width** y_U to every s, t -cut $\delta(U)$.

Definition

A width assignment $\{y_U : \delta(U) \text{ } s, t\text{-cut}\}$ is **feasible** if, for every edge $e \in E$, the **total width** of all cuts containing e is no more than c_e .




Using math: y is feasible if for all e

$$\sum (y_U : \delta(U) \text{ } s, t\text{-cut and } e \in E) \leq c_e$$

Back to the General Case

Consider the **example** on the right with 4 s, t -cuts.



$$U_1 = \{s\}$$

$$U_2 = \{s, a\}$$

$$U_3 = \{s, a, c\}$$

$$U_4 = \{s, a, b, c, d\}$$

Back to the General Case

Consider the **example** on the right with 4 s, t -cuts.

The width assignment


$$y_{U_1} = 3$$

$$y_{U_2} = 1$$

$$y_{U_3} = 2$$

$$y_{U_4} = 1$$

is easily checked to be feasible.



$$U_1 = \{s\}$$

$$U_2 = \{s, a\}$$

$$U_3 = \{s, a, c\}$$


$$U_4 = \{s, a, b, c, d\}$$

Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ } s, t\text{-cut}).$$



$$U_1 = \{s\}$$

$$U_2 = \{s, a\}$$

$$U_3 = \{s, a, c\}$$


$$U_4 = \{s, a, b, c, d\}$$

Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ s,t-cut}).$$



Example:

$$y_{U_1} + y_{U_2} + y_{U_3} + y_{U_4} = 7$$

$$U_1 = \{s\}$$

$$U_2 = \{s, a\}$$

$$U_3 = \{s, a, c\}$$


$$U_4 = \{s, a, b, c, d\}$$

Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ s,t-cut}).$$



Example:

$$y_{U_1} + y_{U_2} + y_{U_3} + y_{U_4} = 7$$

→ Path sa, ac, cb, bt is a **shortest path!**

$$U_1 = \{s\}$$

$$U_2 = \{s, a\}$$

$$U_3 = \{s, a, c\}$$

$$U_4 = \{s, a, b, c, d\}$$

Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ } s, t\text{-cut}).$$

Proof: Consider an s, t -path P . It follows that

$$c(P) = \sum (c_e : e \in P)$$

Example:

$$y_{U_1} + y_{U_2} + y_{U_3} + y_{U_4} = 7$$

→ Path sa, ac, cb, bt is a **shortest path!**



Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ } s, t\text{-cut}).$$

Example:

$$y_{U_1} + y_{U_2} + y_{U_3} + y_{U_4} = 7$$

→ Path sa, ac, cb, bt is a **shortest path!**

Proof: Consider an s, t -path P . It follows that

$$\begin{aligned} c(P) &= \sum (c_e : e \in P) \\ &\geq \sum \left(\sum (y_U : e \in \delta(U)) : e \in P \right) \end{aligned}$$

where the first inequality follows from the feasibility of y .

□

Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ } s, t\text{-cut}).$$

Example:

$$y_{U_1} + y_{U_2} + y_{U_3} + y_{U_4} = 7$$

→ Path sa, ac, cb, bt is a **shortest path!**

Proof: Consider an s, t -path P . It follows that

$$\begin{aligned} c(P) &= \sum (c_e : e \in P) \\ &\geq \sum \left(\sum (y_U : e \in \delta(U)) : e \in P \right) \end{aligned}$$

where the first inequality follows from the feasibility of y .

Note: if $\delta(U)$ is an s, t -cut, then P contains at least one edge from $\delta(U)$.

□

Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ } s, t\text{-cut}).$$

Example:

$$y_{U_1} + y_{U_2} + y_{U_3} + y_{U_4} = 7$$

→ Path sa, ac, cb, bt is a **shortest path!**

Proof: Consider an s, t -path P . It follows that

$$\begin{aligned} c(P) &= \sum (c_e : e \in P) \\ &\geq \sum \left(\sum (y_U : e \in \delta(U)) : e \in P \right) \end{aligned}$$

where the first inequality follows from the feasibility of y .

Note: if $\delta(U)$ is an s, t -cut, then P contains at least one edge from $\delta(U)$.

→ Variable y_U appears **at least** once on the right-hand side above, and hence we obtain the 2nd inequality □

Back to the General Case

Proposition

If y is a **feasible width assignment**, then any s, t -path must have length at least

$$\sum (y_U : U \text{ } s, t\text{-cut}).$$

Example:

$$y_{U_1} + y_{U_2} + y_{U_3} + y_{U_4} = 7$$

→ Path sa, ac, cb, bt is a **shortest path!**

Proof: Consider an s, t -path P . It follows that

$$\begin{aligned} c(P) &= \sum (c_e : e \in P) \\ &\geq \sum \left(\sum (y_U : e \in \delta(U)) : e \in P \right) \\ &\geq \sum (y_U : \delta(U) \text{ } s, t\text{-cut}) \end{aligned}$$


where the first inequality follows from the feasibility of y .

Note: if $\delta(U)$ is an s, t -cut, then P contains at least one edge from $\delta(U)$.

→ Variable y_U appears **at least** once on the right-hand side above, and hence we obtain the 2nd inequality □

One More Example


Question: Can you spot a shortest s, t -path?



One More Example

Question: Can you spot a shortest s, t -path?

→ $P = sa, ac, cd, dt$ of length 7.




One More Example

Question: Can you spot a shortest s, t -path?

→ $P = sa, ac, cd, dt$ of length 7.

Question: Can you prove your guess?




One More Example

Question: Can you spot a shortest s, t -path?

→ $P = sa, ac, cd, dt$ of length 7.

Question: Can you prove your guess?

→ **Yes!** There is a feasible dual width assignment of value 7:



$$y_{\{s\}} = 2$$

$$y_{\{s,a\}} = 1$$

$$y_{\{s,a,c\}} = 1$$

$$y_{\{s,a,c,e\}} = 1$$


$$y_{\{s,a,c,d,e\}} = 1$$

$$y_{\{s,a,b,c,d,e\}} = 1$$

One More Example

Question

(A) In an instance with a shortest path, can we **always** find feasible widths to prove optimality?




One More Example

Question

(A) In an instance with a shortest path, can we **always** find feasible widths to prove optimality?

(B) If so, **how** do we find a path and these widths?



Recap

- A shortest path instance is given by a graph $G = (V, E)$ and non-negative lengths c_e for all $e \in E$.

Recap

- A shortest path instance is given by a graph $G = (V, E)$ and non-negative lengths c_e for all $e \in E$.
- A width assignment $y_U \geq 0$ for all s, t -cuts $\delta(U)$ is **feasible** if

$$\sum (y_U : e \in \delta(U)) \leq c_e$$

for all $e \in E$.

Recap

- A shortest path instance is given by a graph $G = (V, E)$ and non-negative lengths c_e for all $e \in E$.
- A width assignment $y_U \geq 0$ for all s, t -cuts $\delta(U)$ is **feasible** if

$$\sum (y_U : e \in \delta(U)) \leq c_e$$

for all $e \in E$.

- If y is a feasible width assignment and P an s, t -path, then

$$c(P) \geq \sum y_U$$