

## Module 2: Linear Programs (Finding a Feasible Solution)

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

To run Simplex, we need a **feasible** basis.

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

To run Simplex, we need a **feasible** basis.

## Question

How do we find a feasible basis?

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

To run Simplex, we need a **feasible** basis.

## Question

How do we find a feasible basis?

An easier question,

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

To run Simplex, we need a **feasible** basis.

## Question

How do we find a feasible basis?

An easier question,

## Question

How do we find a feasible solution?

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

To run Simplex, we need a **feasible** basis.

## Question

How do we find a feasible basis?

An easier question,

## Question

How do we find a feasible solution?

These two questions are equivalent.

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

To run Simplex, we need a **feasible** basis.

## Question

How do we find a feasible basis?

An easier question,

## Question

How do we find a feasible solution?

These two questions are equivalent.

## Exercise

There is an algorithm that, given a feasible solution, finds a feasible basis.

# The Problem

Consider

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}.$$

To run Simplex, we need a **feasible** basis.

## Question

How do we find a feasible basis?

An easier question,

## Question

How do we find a feasible solution?

These two questions are equivalent.

## Exercise

There is an algorithm that, given a feasible solution, finds a feasible basis.

➡ We will focus on the second question.

# The Key Idea

# The Key Idea

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

# The Key Idea

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

## Algorithm 1

INPUT:  $A, b, c$ , and a **feasible solution**

OUTPUT: Optimal solution/detect LP unbounded.

# The Key Idea

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

## Algorithm 1

INPUT:  $A, b, c$ , and a **feasible solution**

OUTPUT: Optimal solution/detect LP unbounded.

OK  
Simplex +  
exercise

# The Key Idea

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

## Algorithm 1

INPUT:  $A, b, c$ , and a **feasible solution**

OUTPUT: Optimal solution/detect LP unbounded.

OK  
Simplex +  
exercise

## Algorithm 2

INPUT:  $A, b, c$ .

OUTPUT: Feasible solution/detect there is none.

# The Key Idea

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

## Algorithm 1

INPUT:  $A, b, c$ , and a **feasible solution**

OUTPUT: Optimal solution/detect LP unbounded.

OK  
Simplex +  
exercise

## Algorithm 2

INPUT:  $A, b, c$ .

OUTPUT: Feasible solution/detect there is none.

HOW?

# The Key Idea

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

## Algorithm 1

INPUT:  $A, b, c$ , and a **feasible solution**

OUTPUT: Optimal solution/detect LP unbounded.

OK  
Simplex +  
exercise

## Algorithm 2

INPUT:  $A, b, c$ .

OUTPUT: Feasible solution/detect there is none.

HOW?

We will show that...

# The Key Idea

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

## Algorithm 1

INPUT:  $A, b, c$ , and a **feasible solution**

OUTPUT: Optimal solution/detect LP unbounded.

OK  
Simplex +  
exercise

## Algorithm 2

INPUT:  $A, b, c$ .

OUTPUT: Feasible solution/detect there is none.

HOW?

We will show that...

## Proposition

We can use **Algorithm 1** to get **Algorithm 2**.

# A First Example

# A First Example

Problem: Find a feasible solution/detect none exist for

$$\max (1, 2, -1, 3)x$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ -2 & -9 & 0 & 3 \end{pmatrix} x = \begin{pmatrix} 7 \\ -13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

# A First Example

Problem: Find a feasible solution/detect none exist for

$$\max (1, 2, -1, 3)x$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ -2 & -9 & 0 & 3 \end{pmatrix} x = \begin{pmatrix} 7 \\ -13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

## Remark

It does not depend on the objective function.

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ -2 & -9 & 0 & 3 \end{pmatrix} x = \begin{pmatrix} 7 \\ -13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ -2 & -9 & 0 & 3 \end{pmatrix} x = \begin{pmatrix} 7 \\ -13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative.

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ -2 & -9 & 0 & 3 \end{pmatrix} x = \begin{pmatrix} 7 \\ -13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative.

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

## Remark

The auxiliary problem is

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (*)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

## Remark

The auxiliary problem is

- feasible, since  $(0, 0, 0, 0, 7, 13)^\top$  is a solution, and

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (*)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

## Remark

The auxiliary problem is

- feasible, since  $(0, 0, 0, 0, 7, 13)^\top$  is a solution, and
- bounded, as 0 is the lower bound.

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (*)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

## Remark

The auxiliary problem is

- feasible, since  $(0, 0, 0, 0, 7, 13)^\top$  is a solution, and
- bounded, as 0 is the lower bound.

➡ Therefore, the auxiliary problem has an optimal solution.

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (*)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(2, 1, 0, 0, 0, 0)^\top$  is an optimal solution to the auxiliary problem,

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (*)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(2, 1, 0, 0, 0, 0)^\top$  is an optimal solution to the auxiliary problem,

since  $x_5 = x_6 = 0$ .

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 5 & 2 & 1 \\ 2 & 9 & 0 & -3 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad x_5 + x_6$$

s.t.

$$\begin{pmatrix} 1 & 5 & 2 & 1 & 1 & 0 \\ 2 & 9 & 0 & -3 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 7 \\ 13 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_5, x_6$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(2, 1, 0, 0, 0, 0)^\top$  is an optimal solution to the auxiliary problem,

since  $x_5 = x_6 = 0$ .

➡ Therefore,  $(2, 1, 0, 0)^\top$  is a feasible solution to  $(\star)$ .

## A Second Example

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative.

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_4, x_5$  are the  
**auxiliary variables**

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_4, x_5$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_4, x_5$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(0, 0, 1, 0, 3)^\top$  is an optimal solution to the auxiliary problem.

## A Second Example

Problem: Find a feasible solution/detect none exist for

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$x_4, x_5$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(0, 0, 1, 0, 3)^\top$  is an optimal solution to the auxiliary problem.

However,  $(0, 0, 1)^\top$  is **NOT** a solution to  $(\star)$ .

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution  $(0, 0, 1, 0, 3)^\top$

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution  $(0, 0, 1, 0, 3)^\top$

optimal value  $= 0 + 3 = 3$ .

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution  $(0, 0, 1, 0, 3)^\top$

optimal value  $= 0 + 3 = 3$ .

## Claim

$(\star)$  does not have a solution.

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution  $(0, 0, 1, 0, 3)^\top$

optimal value  $= 0 + 3 = 3$ .

## Claim

$(\star)$  does not have a solution.

## Proof

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution  $(0, 0, 1, 0, 3)^\top$

optimal value  $= 0 + 3 = 3$ .

## Claim

$(\star)$  does not have a solution.

## Proof

Suppose, for a contradiction,  $(\star)$  has a solution  $x'_1, x'_2, x'_3$ .

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution  $(0, 0, 1, 0, 3)^\top$

optimal value  $= 0 + 3 = 3$ .

## Claim

$(\star)$  does not have a solution.

## Proof

Suppose, for a contradiction,  $(\star)$  has a solution  $x'_1, x'_2, x'_3$ .

Then,  $(x'_1, x'_2, x'_3, 0, 0)^\top$  is a feasible solution to the auxiliary problem,

$$\begin{pmatrix} 5 & 1 & 1 \\ -1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 5 & 1 & 1 & 1 & 0 \\ -1 & 1 & 2 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution  $(0, 0, 1, 0, 3)^\top$

optimal value  $= 0 + 3 = 3$ .

## Claim

$(\star)$  does not have a solution.

## Proof

Suppose, for a contradiction,  $(\star)$  has a solution  $x'_1, x'_2, x'_3$ .

Then,  $(x'_1, x'_2, x'_3, 0, 0)^\top$  is a feasible solution to the auxiliary problem,

but that solution has of value 0. This is a contradiction.

# Formalize

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$\begin{aligned} \min \quad & z = x_{n+1} + \dots + x_{n+m} \\ \text{s.t.} \quad & \left( \begin{array}{c c} A & I \end{array} \right) x = b \\ & x \geq \mathbf{0} \end{aligned}$
--

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (*)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$\begin{aligned} \min \quad & z = x_{n+1} + \dots + x_{n+m} \\ \text{s.t.} \quad & \left( \begin{array}{c c} A & I \end{array} \right) x = b \\ & x \geq \mathbf{0} \end{aligned}$
--

$x_{n+1}, \dots, x_{n+m}$  are the  
**auxiliary variables**

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$
$$x \geq \mathbf{0}$$

$x_{n+1}, \dots, x_{n+m}$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$
$$x \geq \mathbf{0}$$

$x_{n+1}, \dots, x_{n+m}$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$  is an optimal solution to the auxiliary problem.

# Formalize

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$\begin{aligned} \min \quad & z = x_{n+1} + \dots + x_{n+m} \\ \text{s.t.} \quad & \left( \begin{array}{c c} A & I \end{array} \right) x = b \\ & x \geq \mathbf{0} \end{aligned}$
--

$x_{n+1}, \dots, x_{n+m}$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$  is an optimal solution to the auxiliary problem.

## Proposition

If  $z = 0$ , then  $(x_1, \dots, x_n)^\top$  is a solution to  $(\star)$ .

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$

$$x \geq \mathbf{0}$$

$x_{n+1}, \dots, x_{n+m}$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$  is an optimal solution to the auxiliary problem.

## Proposition

If  $z = 0$ , then  $(x_1, \dots, x_n)^\top$  is a solution to  $(\star)$ .

## Proof

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$

$$x \geq \mathbf{0}$$

$x_{n+1}, \dots, x_{n+m}$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$  is an optimal solution to the auxiliary problem.

## Proposition

If  $z = 0$ , then  $(x_1, \dots, x_n)^\top$  is a solution to  $(\star)$ .

## Proof

When  $z = 0$ , we must have  $x_{n+1} = \dots = x_{n+m} = 0$ .

Problem: Find a feasible solution/detect none exist for

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that  $b$  is non-negative.

**Step 2.** Construct the **auxiliary problem** ( $A$   $m \times n$  matrix).

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$

$$x \geq \mathbf{0}$$

$x_{n+1}, \dots, x_{n+m}$  are the  
**auxiliary variables**

**Step 3.** Solve the **auxiliary problem** using **Algorithm 1**.

$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^T$  is an optimal solution to the auxiliary problem.

## Proposition

When  $z > 0$ , then  $(\star)$  has no solution.

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$
$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution

$$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$$

## Proposition

When  $z > 0$ , then  $(\star)$  has no solution.

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$
$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution

$$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$$

## Proposition

When  $z > 0$ , then  $(\star)$  has no solution.

## Proof

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$
$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution

$$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$$

## Proposition

When  $z > 0$ , then  $(\star)$  has no solution.

## Proof

Suppose, for a contradiction,  $(\star)$  has a solution  $x'_1, \dots, x'_n$ .

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\begin{array}{ll} \min & z = x_{n+1} + \dots + x_{n+m} \\ \text{s.t.} & \\ & \left( \begin{array}{c|c} A & I \end{array} \right) x = b \\ & x \geq \mathbf{0} \end{array}$$

the auxiliary problem

optimal solution

$$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$$

## Proposition

When  $z > 0$ , then  $(\star)$  has no solution.

## Proof

Suppose, for a contradiction,  $(\star)$  has a solution  $x'_1, \dots, x'_n$ .

Then  $(x'_1, \dots, x'_n, \mathbf{0}, \dots, \mathbf{0})^\top$  is a feasible solution to the auxiliary problem,

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\min \quad z = x_{n+1} + \dots + x_{n+m}$$

s.t.

$$\left( \begin{array}{c|c} A & I \end{array} \right) x = b$$
$$x \geq \mathbf{0}$$

the auxiliary problem

optimal solution

$$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$$

## Proposition

When  $z > 0$ , then  $(\star)$  has no solution.

## Proof

Suppose, for a contradiction,  $(\star)$  has a solution  $x'_1, \dots, x'_n$ .

Then  $(x'_1, \dots, x'_n, 0, \dots, 0)^\top$  is a feasible solution to the auxiliary problem,

but that solution has of value 0.

$$Ax = b \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

$$\begin{array}{ll} \min & z = x_{n+1} + \dots + x_{n+m} \\ \text{s.t.} & \\ & \left( \begin{array}{c|c} A & I \end{array} \right) x = b \\ & x \geq \mathbf{0} \end{array}$$

the auxiliary problem

optimal solution

$$(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m})^\top$$

## Proposition

When  $z > 0$ , then  $(\star)$  has no solution.

## Proof

Suppose, for a contradiction,  $(\star)$  has a solution  $x'_1, \dots, x'_n$ .

Then  $(x'_1, \dots, x'_n, \mathbf{0}, \dots, \mathbf{0})^\top$  is a feasible solution to the auxiliary problem, but that solution has of value 0. This is a contradiction.

# The 2-Phase Method

# The 2-Phase Method

To solve

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\},$$

# The 2-Phase Method

To solve

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\},$$

we proceed in two phases:

# The 2-Phase Method

To solve

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\},$$

we proceed in two phases:

**Phase 1.** Find a feasible solution/detect none exist.

# The 2-Phase Method

To solve

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\},$$

we proceed in two phases:

**Phase 1.** Find a feasible solution/detect none exist.

**Phase 2.** Given that feasible solution:  
find an optimal solution/detect LP unbounded.

# The 2-Phase Method

To solve

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\},$$

we proceed in two phases:

**Phase 1.** Find a feasible solution/detect none exist.

**Phase 2.** Given that feasible solution:  
find an optimal solution/detect LP unbounded.

## Example

Solve the following LP,

$$\begin{array}{ll} \max & (1, 1, 1)x \\ \text{s.t.} & \\ & \begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \\ & x \geq \mathbf{0} \end{array}$$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative.

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. **OK**

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\min \quad z = x_4 + x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

NOT in SEF

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

To rewrite  $B = \{4, 5\}$  in canonical form, you can

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

To rewrite  $B = \{4, 5\}$  in canonical form, you can

- use the formulae, OR

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

To rewrite  $B = \{4, 5\}$  in canonical form, you can

- use the formulae, OR
- notice  $A_B = I$  and rewrite the objective function as follows...

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

$$z = (0 \quad 0 \quad 0 \quad -1 \quad -1)x$$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

$$z = (0 \quad 0 \quad 0 \quad -1 \quad -1)x$$

$$0 = (1 \quad 2 \quad -1 \quad 1 \quad 0)x - 4$$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the auxiliary problem.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

$$z = (0 \quad 0 \quad 0 \quad -1 \quad -1)x$$

$$0 = (1 \quad 2 \quad -1 \quad 1 \quad 0)x - 4$$

$$0 = (1 \quad -1 \quad 1 \quad 0 \quad 1)x - 4$$

---

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the auxiliary problem.

$$\max \quad z = -x_4 - x_5$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

NOT in canonical form

$$z = (0 \quad 0 \quad 0 \quad -1 \quad -1)x$$

$$0 = (1 \quad 2 \quad -1 \quad 1 \quad 0)x - 4$$

$$0 = (1 \quad -1 \quad 1 \quad 0 \quad 1)x - 4$$

---

$$z = (2 \quad 1 \quad 0 \quad 0 \quad 0)x - 8$$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max z = (2 \ 1 \ 0 \ 0 \ 0) - 8$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max z = (2 \ 1 \ 0 \ 0 \ 0) - 8$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

**canonical form for  $B$**

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (*)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max z = (2 \ 1 \ 0 \ 0 \ 0) - 8$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

**canonical form for  $B$**

**Step 3.** Solve the **auxiliary problem** using **Simplex**, starting from  $B$ .

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max z = (2 \ 1 \ 0 \ 0 \ 0) - 8$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

**canonical form for  $B$**

**Step 3.** Solve the **auxiliary problem** using **Simplex**, starting from  $B$ .

$B = \{1, 4\}$  is an optimal basis with the basic solution  $(4, 0, 0, 0, 0)^\top$ .

**Phase 1.** Find a feasible solution/detect none exist for

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad \text{and} \quad x \geq \mathbf{0} \quad (\star)$$

**Step 1.** Multiply the equations such that the RHS is non-negative. OK

**Step 2.** Construct the **auxiliary problem**.

$$\max z = (2 \ 1 \ 0 \ 0 \ 0) - 8$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 & 1 & 0 \\ 1 & -1 & 1 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

In SEF

feasible basis  $B = \{4, 5\}$

**canonical form for  $B$**

**Step 3.** Solve the **auxiliary problem** using **Simplex**, starting from  $B$ .

$B = \{1, 4\}$  is an optimal basis with the basic solution  $(4, 0, 0, 0, 0)^\top$ .

$z = 0$  implies that  $(4, 0, 0)^\top$  is a feasible solution for  $(\star)$ .

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\max (2, -1, 2)x$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\max (2, -1, 2)x$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

## Remark

$(4, 0, 0)^\top$  is a **basic** solution.

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\max (2, -1, 2)x$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

## Remark

$(4, 0, 0)^\top$  is a **basic** solution.

## Exercise

Show that this will always be the case!

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\max (2, -1, 2)x$$

s.t.

$$\underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\max (2, -1, 2)x$$

s.t.

$$\underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

## Question

For what basis,  $B$ , is  $x = (4, 0, 0)^\top$  a basic solution?

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\max (2, -1, 2)x$$

s.t.

$$\underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

## Question

For what basis,  $B$ , is  $x = (4, 0, 0)^\top$  a basic solution?

$$x_1 \neq 0$$

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\max (2, -1, 2)x$$

s.t.

$$\underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

## Question

For what basis,  $B$ , is  $x = (4, 0, 0)^\top$  a basic solution?

$$x_1 \neq 0 \implies 1 \in B$$

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\begin{array}{l} \max \quad (2, -1, 2)x \\ \text{s.t.} \\ \underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix} \\ x \geq \mathbf{0} \end{array}$$

## Question

For what basis,  $B$ , is  $x = (4, 0, 0)^\top$  a basic solution?

$$x_1 \neq 0 \implies 1 \in B$$

Cardinality of maximal set of independent columns of  $A =$

Cardinality of maximal set of independent rows of  $A = 2$

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\begin{array}{ll} \max & (2, -1, 2)x \\ \text{s.t.} & \\ & \underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix} \\ & x \geq \mathbf{0} \end{array}$$

## Question

For what basis,  $B$ , is  $x = (4, 0, 0)^\top$  a basic solution?

$$x_1 \neq 0 \implies 1 \in B$$

Cardinality of maximal set of independent columns of  $A =$   
Cardinality of maximal set of independent rows of  $A = 2$

Thus, for some  $i \in \{2, 3\}$ , columns 1 and  $i$  of  $A$  are independent.

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\begin{array}{l} \max \quad (2, -1, 2)x \\ \text{s.t.} \\ \underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix} \\ x \geq \mathbf{0} \end{array}$$

## Question

For what basis,  $B$ , is  $x = (4, 0, 0)^\top$  a basic solution?

$$x_1 \neq 0 \implies 1 \in B$$

Cardinality of maximal set of independent columns of  $A =$   
Cardinality of maximal set of independent rows of  $A = 2$

Thus, for some  $i \in \{2, 3\}$ , columns 1 and  $i$  of  $A$  are independent.

In this case, we can pick  $i = 2$ .

**Phase 1.**  $(4, 0, 0)^\top$  is a feasible solution for

$$\begin{array}{l} \max \quad (2, -1, 2)x \\ \text{s.t.} \\ \underbrace{\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix}}_A x = \begin{pmatrix} 1 \\ 3 \end{pmatrix} \\ x \geq \mathbf{0} \end{array}$$

## Question

For what basis,  $B$ , is  $x = (4, 0, 0)^\top$  a basic solution?

$$x_1 \neq 0 \implies 1 \in B$$

Cardinality of maximal set of independent columns of  $A =$   
Cardinality of maximal set of independent rows of  $A = 2$

Thus, for some  $i \in \{2, 3\}$ , columns 1 and  $i$  of  $A$  are independent.

In this case, we can pick  $i = 2$ . In particular,  $B = \{1, 2\}$  is a basis.

**Phase 2.** Find an optimal solution/detect LP unbounded.

$$\max (1, 1, 1)x$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$B = \{1, 2\}$  is a feasible basis (from [Phase 1](#)).

**Phase 2.** Find an optimal solution/detect LP unbounded.

$$\max (1, 1, 1)x$$

s.t.

$$\begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$x \geq \mathbf{0}$$

$B = \{1, 2\}$  is a feasible basis (from [Phase 1](#)).

We can now solve the problem using Simplex, starting from  $B$ .

**Phase 2.** Find an optimal solution/detect LP unbounded.

$$\begin{array}{ll} \max & (1, 1, 1)x \\ \text{s.t.} & \\ & \begin{pmatrix} 1 & 2 & -1 \\ 1 & -1 & 1 \end{pmatrix} x = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \\ & x \geq \mathbf{0} \end{array}$$

$B = \{1, 2\}$  is a feasible basis (from **Phase 1**).

We can now solve the problem using Simplex, starting from  $B$ .



$x = (0, 8, 12)^\top$  is an optimal solution.

# Consequences

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

(A) it is infeasible,

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

## Proof

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

## Proof

Run 2-Phase method with Simplex using Bland's rule.

# Consequences

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

## Proof

Run 2-Phase method with Simplex using Bland's rule.  
(Recall that Bland's rule ensures that Simplex terminates.)

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

## Remark

A finite number of basis implies a finite number of basic solutions.

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$

Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

## Remark

A finite number of basis implies a finite number of basic solutions.

If a LP has at least two feasible solutions and one optimal solution,

## Theorem

$$\max \{c^\top x : Ax = b, x \geq \mathbf{0}\}$$


Exactly one of the following holds for the LP:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution that is **basic**.

## Remark

A finite number of basis implies a finite number of basic solutions.

If a LP has at least two feasible solutions and one optimal solution,



# Fundamental Theorem of Linear Programming

# Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP.

## Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP. Exactly one of the following holds:

## Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP. Exactly one of the following holds:

(A) it is infeasible,

## Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP. Exactly one of the following holds:

- (A) it is infeasible,
- (B) it is unbounded, or

## Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP. Exactly one of the following holds:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution.

## Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP. Exactly one of the following holds:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution.

## Proof

## Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP. Exactly one of the following holds:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution.

### Proof

Convert the LP into an **equivalent** LP in SEF.

## Fundamental Theorem of Linear Programming

Consider an **arbitrary** LP. Exactly one of the following holds:

- (A) it is infeasible,
- (B) it is unbounded, or
- (C) it has an optimal solution.

### Proof

Convert the LP into an **equivalent** LP in SEF.

Apply the previous theorem.

## Recap

## Recap

- Using 2-Phase + the Simplex algorithm, we can solve arbitrary LPs.

## Recap

- Using 2-Phase + the Simplex algorithm, we can solve arbitrary LPs.
- We proved the fundamental theorem of linear programming.

## Recap

- Using 2-Phase + the Simplex algorithm, we can solve arbitrary LPs.
- We proved the fundamental theorem of linear programming.

We have given a bare bone version of the Simplex procedure.

## Recap

- Using 2-Phase + the Simplex algorithm, we can solve arbitrary LPs.
- We proved the fundamental theorem of linear programming.

We have given a bare bone version of the Simplex procedure.

**Careful implementation** is key to having a practical algorithm.

## Recap

- Using 2-Phase + the Simplex algorithm, we can solve arbitrary LPs.
- We proved the fundamental theorem of linear programming.

We have given a bare bone version of the Simplex procedure.

**Careful implementation** is key to having a practical algorithm.

If Simplex were a bike,

## Recap

- Using 2-Phase + the Simplex algorithm, we can solve arbitrary LPs.
- We proved the fundamental theorem of linear programming.

We have given a bare bone version of the Simplex procedure.

**Careful implementation** is key to having a practical algorithm.

If Simplex were a bike,



Sergio Schnitzler/Hemera/Thinkstock

State of the art implementation

## Recap

- Using 2-Phase + the Simplex algorithm, we can solve arbitrary LPs.
- We proved the fundamental theorem of linear programming.

We have given a bare bone version of the Simplex procedure.

Careful implementation is key to having a practical algorithm.

If Simplex were a bike,



Sergio Schnitzler/Hemera/Thinkstock

State of the art implementation



gamegfx/iStock/Thinkstock

Our implementation