

## **Module 6: Nonlinear Programs (Convexity)**

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\min f(x)$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

where

$$f : \mathbb{R}^n \rightarrow \mathbb{R}, \text{ and}$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

where

$$f : \mathbb{R}^n \rightarrow \mathbb{R}, \text{ and}$$

$$g_i : \mathbb{R}^n \rightarrow \mathbb{R} \text{ for } i = 1, \dots, k.$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

where

$$f : \mathbb{R}^n \rightarrow \mathbb{R}, \text{ and}$$

$$g_i : \mathbb{R}^n \rightarrow \mathbb{R} \text{ for } i = 1, \dots, k.$$

## Remark

There aren't any restrictions regarding the type of functions.

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

where

$$f : \mathbb{R}^n \rightarrow \mathbb{R}, \text{ and}$$

$$g_i : \mathbb{R}^n \rightarrow \mathbb{R} \text{ for } i = 1, \dots, k.$$

## Remark

There aren't any restrictions regarding the type of functions.

This is a very general model, but NLPs can be very hard to solve!

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

$$\begin{array}{ll} \min & x_2 \\ \text{s.t.} & \\ & -x_1^2 - x_2 + 2 \leq 0 \\ & x_2 - \frac{3}{2} \leq 0 \\ & x_1 - \frac{3}{2} \leq 0 \\ & -x_1 - 2 \leq 0 \end{array}$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

$$\begin{array}{ll} \min & \underbrace{x_2}_{f(x)} \\ \text{s.t.} & \\ & -x_1^2 - x_2 + 2 \leq 0 \\ & x_2 - \frac{3}{2} \leq 0 \\ & x_1 - \frac{3}{2} \leq 0 \\ & -x_1 - 2 \leq 0 \end{array}$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

$$\min \quad x_2$$

s.t.

$$\underbrace{-x_1^2 - x_2 + 2}_{g_1(x)} \leq 0$$

$$x_2 - \frac{3}{2} \leq 0$$

$$x_1 - \frac{3}{2} \leq 0$$

$$-x_1 - 2 \leq 0$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

$$\min \quad x_2$$

s.t.

$$-x_1^2 - x_2 + 2 \leq 0$$

$$\underbrace{x_2 - \frac{3}{2}}_{g_2(x)} \leq 0$$

$$x_1 - \frac{3}{2} \leq 0$$

$$-x_1 - 2 \leq 0$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

$$\begin{array}{ll} \min & x_2 \\ \text{s.t.} & \\ & -x_1^2 - x_2 + 2 \leq 0 \\ & x_2 - \frac{3}{2} \leq 0 \\ & \underbrace{x_1 - \frac{3}{2}}_{g_3(x)} \leq 0 \\ & -x_1 - 2 \leq 0 \end{array}$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

$$\begin{array}{ll} \min & x_2 \\ \text{s.t.} & \\ & -x_1^2 - x_2 + 2 \leq 0 \\ & x_2 - \frac{3}{2} \leq 0 \\ & x_1 - \frac{3}{2} \leq 0 \\ & \underbrace{-x_1 - 2}_{g_4(x)} \leq 0 \end{array}$$

min  $x_2$

s.t.


$$-x_1^2 - x_2 + 2 \leq 0$$

$$x_2 - 3/2 \leq 0$$

$$x_1 - 3/2 \leq 0$$

$$-x_1 - 2 \leq 0$$

(1)  $x_2 \geq 2 - x_1^2$ .



min  $x_2$

s.t.

$$-x_1^2 - x_2 + 2 \leq 0$$


$$x_2 - 3/2 \leq 0$$

$$x_1 - 3/2 \leq 0$$

$$-x_1 - 2 \leq 0$$

(1)  $x_2 \geq 2 - x_1^2$ .

(2)  $x_2 \leq \frac{3}{2}$ .



min  $x_2$

s.t.

$$-x_1^2 - x_2 + 2 \leq 0$$

$$x_2 - 3/2 \leq 0$$


$$x_1 - 3/2 \leq 0$$

$$-x_1 - 2 \leq 0$$

(1)  $x_2 \geq 2 - x_1^2$ .

(2)  $x_2 \leq \frac{3}{2}$ .

(3)  $x_1 \leq \frac{3}{2}$ .



$$\min x_2$$

s.t.

$$-x_1^2 - x_2 + 2 \leq 0$$

$$x_2 - 3/2 \leq 0$$

$$x_1 - 3/2 \leq 0$$


$$-x_1 - 2 \leq 0$$

$$(1) x_2 \geq 2 - x_1^2.$$

$$(2) x_2 \leq \frac{3}{2}.$$

$$(3) x_1 \leq \frac{3}{2}.$$

$$(4) x_1 \geq -2.$$



min  $x_2$


s.t.

$$-x_1^2 - x_2 + 2 \leq 0$$

$$x_2 - 3/2 \leq 0$$

$$x_1 - 3/2 \leq 0$$

$$-x_1 - 2 \leq 0$$



FEASIBLE REGION

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\min f(x)$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

(P)

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

## Remark

We may assume  $f(x)$  is a **linear function**, i.e.,  $f(x) = c^\top x$ .

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

## Remark

We may assume  $f(x)$  is a **linear function**, i.e.,  $f(x) = c^\top x$ .

We can rewrite (P) as

$$\begin{array}{ll} \min & \lambda \\ \text{s.t.} & \\ & \lambda \geq f(x) \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{Q})$$

## Definition

A **Nonlinear Program** (NLP) is a problem of the form:

$$\begin{array}{ll} \min & f(x) \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

## Remark

We may assume  $f(x)$  is a **linear function**, i.e.,  $f(x) = c^\top x$ .

We can rewrite (P) as

$$\begin{array}{ll} \min & \lambda \\ \text{s.t.} & \\ & \lambda \geq f(x) \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{Q})$$

The optimal solution to (Q) will have  $\lambda = f(x)$ .

# Nonlinear Programs Generalize Linear Programs

$$\max \quad x_1 + x_2$$

s.t.

$$2x_1 - x_2 \geq 3$$

$$x_1 - x_2 = 4$$

$$x_1, x_2 \geq 0$$

# Nonlinear Programs Generalize Linear Programs

$$\max \quad x_1 + x_2$$

s.t.

$$2x_1 - x_2 \geq 3$$

$$x_1 - x_2 = 4$$

$$x_1, x_2 \geq 0$$

$$\min \quad -x_1 - x_2$$

s.t.

$$-2x_1 + x_2 + 3 \leq 0$$

$$x_1 - x_2 - 4 \leq 0$$

$$-x_1 + x_2 + 4 \leq 0$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0$$

# Nonlinear Programs Generalize Linear Programs

$$\max \quad x_1 + x_2$$

s.t.

$$2x_1 - x_2 \geq 3$$

$$x_1 - x_2 = 4$$

$$x_1, x_2 \geq 0$$

$$\min \quad -x_1 - x_2$$

s.t.

$$-2x_1 + x_2 + 3 \leq 0$$

$$x_1 - x_2 - 4 \leq 0$$

$$-x_1 + x_2 + 4 \leq 0$$

$$-x_1 \leq 0$$

$$-x_2 \leq 0$$

Nonlinear Programs can also generalize **INTEGER PROGRAMS!**

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \in \{0, 1\} \quad (j = 1, \dots, n)$$

0, 1 IP

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \in \{0, 1\} \quad (j = 1, \dots, n)$$

0, 1 IP

## Idea

$$x_j \in \{0, 1\}$$

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \in \{0, 1\} \quad (j = 1, \dots, n)$$

0, 1 IP

## Idea

$$x_j \in \{0, 1\} \quad \iff \quad x_j(1 - x_j) = 0$$

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \in \{0, 1\} \quad (j = 1, \dots, n)$$

0,1 IP

## Idea

$$x_j \in \{0, 1\} \quad \iff \quad x_j(1 - x_j) = 0$$

$$\min \quad -c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j(1 - x_j) \leq 0 \quad (j = 1, \dots, n)$$

$$-x_j(1 - x_j) \leq 0 \quad (j = 1, \dots, n)$$

Quadratic NLP

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \in \{0, 1\} \quad (j = 1, \dots, n)$$

0, 1 IP

## Idea

$$x_j \in \{0, 1\} \quad \iff \quad x_j(1 - x_j) = 0$$

$$\min \quad -c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j(1 - x_j) \leq 0 \quad (j = 1, \dots, n)$$

$$-x_j(1 - x_j) \leq 0 \quad (j = 1, \dots, n)$$

Quadratic NLP

## Remark

0, 1 IPs are hard to solve; thus, quadratic NLPs are also hard to solve.

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \text{ integer} \quad (j = 1, \dots, n)$$

pure IP

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \text{ integer} \quad (j = 1, \dots, n)$$

pure IP

## Idea

$x_j$  integer

# Nonlinear Programs Generalize **Integer** Programs

$$\max c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \text{ integer } (j = 1, \dots, n)$$

pure IP

## Idea

$$x_j \text{ integer} \iff \sin(\pi x_j) = 0.$$

# Nonlinear Programs Generalize Integer Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \text{ integer} \quad (j = 1, \dots, n)$$

pure IP

## Idea

$$x_j \text{ integer} \quad \iff \quad \sin(\pi x_j) = 0.$$

$$\min \quad -c^\top x$$

s.t.

$$Ax \leq b$$

$$\sin(\pi x_j) = 0 \quad (j = 1, \dots, n)$$

# Nonlinear Programs Generalize **Integer** Programs

$$\max \quad c^\top x$$

s.t.

$$Ax \leq b$$

$$x_j \text{ integer} \quad (j = 1, \dots, n)$$

pure IP

## Idea

$$x_j \text{ integer} \quad \iff \quad \sin(\pi x_j) = 0.$$

$$\min \quad -c^\top x$$

s.t.

$$Ax \leq b$$

$$\sin(\pi x_j) = 0 \quad (j = 1, \dots, n)$$

## Remark

IPs are hard to solve; thus, NLPs are also hard to solve.

## Question

What makes solving an NLP hard?

## Question

What makes solving an NLP hard?

META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

## Question

What makes solving an NLP hard?

META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .

## Question

What makes solving an NLP hard?

META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.

## Question

What makes solving an NLP hard?

META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.

## Question

What makes solving an NLP hard?

META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.



## META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.



## META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.



min  $x_2$

s.t.

$$-x_1^2 - x_2 + 2 \leq 0$$

$$x_2 - 3/2 \leq 0$$

$$x_1 - 3/2 \leq 0$$

$$-x_1 - 2 \leq 0$$

# META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.



$$\min x_2$$


s.t.

$$-x_1^2 - x_2 + 2 \leq 0$$

$$x_2 - 3/2 \leq 0$$

$$x_1 - 3/2 \leq 0$$

$$-x_1 - 2 \leq 0$$




## META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.



$a$  is an optimal solution




## META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM

- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.



$a$  is an optimal solution

no better solution around  $b$



## META STRATEGY FOR SOLVING AN OPTIMIZATION PROBLEM


- Find a feasible solution  $x$ .
- If  $x$  is optimal, STOP.
- Starting with  $x$ , find a “better” feasible solution.



$a$  is an optimal solution

no better solution around  $b$

$b$  is a local optimum



## Definition

Consider

$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

## Definition

Consider

$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum**

## Definition

Consider

$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum** if there exists  $\delta > 0$  such that

## Definition

Consider

$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum** if there exists  $\delta > 0$  such that

$$\forall x' \in S \quad \text{where} \quad \|x' - x\| \leq \delta \quad \text{we have} \quad f(x) \leq f(x').$$

## Definition

Consider


$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum** if there exists  $\delta > 0$  such that

$$\forall x' \in S \quad \text{where} \quad \|x' - x\| \leq \delta \quad \text{we have} \quad f(x) \leq f(x').$$

$$\min\{x_2 : x \in S\}$$

$b$  is a **local optimum**



## Definition

Consider


$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum** if there exists  $\delta > 0$  such that

$$\forall x' \in S \quad \text{where} \quad \|x' - x\| \leq \delta \quad \text{we have} \quad f(x) \leq f(x').$$

$$\min\{x_2 : x \in S\}$$

$b$  is a **local optimum**



# Convexity Helps

## Definition

Consider

$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum** if there exists  $\delta > 0$  such that

$$\forall x' \in S \quad \text{where} \quad \|x' - x\| \leq \delta \quad \text{we have} \quad f(x) \leq f(x').$$

# Convexity Helps

## Definition

Consider

$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum** if there exists  $\delta > 0$  such that

$$\forall x' \in S \quad \text{where} \quad \|x' - x\| \leq \delta \quad \text{we have} \quad f(x) \leq f(x').$$

## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

# Convexity Helps

## Definition

Consider

$$\min \{f(x) : x \in S\}. \quad (\text{P})$$

$x \in S$  is a **local optimum** if there exists  $\delta > 0$  such that

$$\forall x' \in S \quad \text{where} \quad \|x' - x\| \leq \delta \quad \text{we have} \quad f(x) \leq f(x').$$

## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.


## Proposition

Consider

$$\min \{c^T x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

Proof




## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

Proof



## Proposition


Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof

Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .



## Proposition

Consider


$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof

Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$


If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof

Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.


## Proof

Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .

Since  $S$  is convex,  $y \in S$ .



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof


Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .

Since  $S$  is convex,  $y \in S$ .

$$c^\top y$$



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof


Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .

Since  $S$  is convex,  $y \in S$ .

$$c^\top y = c^\top (\lambda x' + (1 - \lambda)x)$$



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof


Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .

Since  $S$  is convex,  $y \in S$ .

$$\begin{aligned} c^\top y &= c^\top (\lambda x' + (1 - \lambda)x) \\ &= \underbrace{\lambda}_{>0} \underbrace{c^\top x'}_{< c^\top x} + (1 - \lambda)c^\top x \end{aligned}$$



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof


Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .

Since  $S$  is convex,  $y \in S$ .

$$\begin{aligned} c^\top y &= c^\top (\lambda x' + (1 - \lambda)x) \\ &= \underbrace{\lambda}_{>0} \underbrace{c^\top x'}_{< c^\top x} + (1 - \lambda)c^\top x \\ &< \lambda c^\top x + (1 - \lambda)c^\top x \end{aligned}$$



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof


Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .

Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .

Since  $S$  is convex,  $y \in S$ .

$$\begin{aligned} c^\top y &= c^\top (\lambda x' + (1 - \lambda)x) \\ &= \underbrace{\lambda}_{>0} \underbrace{c^\top x'}_{< c^\top x} + (1 - \lambda)c^\top x \\ &< \lambda c^\top x + (1 - \lambda)c^\top x \\ &= c^\top x \end{aligned}$$



## Proposition

Consider

$$\min \{c^\top x : x \in S\}. \quad (\text{P})$$

If  $S$  is **convex** and  $x$  is a **local optimum**, then  $x$  is optimal.

## Proof

Suppose  $\exists x' \in S$  with  $c^\top x' < c^\top x$ .


Let  $y = \lambda x' + (1 - \lambda)x$  for  $\lambda > 0$  small.

As  $\lambda$  small  $\|y - x\| \leq \delta$ .

Since  $S$  is convex,  $y \in S$ .

$$\begin{aligned} c^\top y &= c^\top (\lambda x' + (1 - \lambda)x) \\ &= \underbrace{\lambda}_{>0} \underbrace{c^\top x'}_{< c^\top x} + (1 - \lambda)c^\top x \\ &< \lambda c^\top x + (1 - \lambda)c^\top x \\ &= c^\top x \end{aligned}$$

A contradiction.



Consider

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

(P)

Consider

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

Goal: Study a case where the feasible region of (P) is **convex**.

Consider

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

Goal: Study a case where the feasible region of (P) is **convex**.

- We will define **convex functions**

Consider

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

Goal: Study a case where the feasible region of (P) is **convex**.

- We will define **convex functions**
- We will prove

## Proposition

If  $g_1, \dots, g_k$  are all convex, then the feasible region of (P) is convex.

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex**

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b)$$

## Definition


Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$




CONVEX FUNCTION!

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$




CONVEX FUNCTION!

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$



CONVEX FUNCTION!

## Definition


Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,


$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$



## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$



NOT A CONVEX FUNCTION!

## Definition


Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$

## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,


$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$



## Definition

Function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if for all  $a, b \in \mathbb{R}^n$ ,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \quad \text{for all } 0 \leq \lambda \leq 1.$$



CONVEX FUNCTION!

## Example

We claim that  $f(x) = x^2$  is convex.

## Example

We claim that  $f(x) = x^2$  is convex.

Pick  $a, b \in \mathfrak{R}$  and pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

## Example

We claim that  $f(x) = x^2$  is convex.

Pick  $a, b \in \mathfrak{R}$  and pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To check:

$$[\lambda a + (1 - \lambda)b]^2 \stackrel{?}{\leq} \lambda a^2 + (1 - \lambda)b^2.$$

## Example

We claim that  $f(x) = x^2$  is convex.

Pick  $a, b \in \mathfrak{R}$  and pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To check:

$$[\lambda a + (1 - \lambda)b]^2 \stackrel{?}{\leq} \lambda a^2 + (1 - \lambda)b^2.$$

We may assume that  $\lambda \neq 0, 1$ .

## Example

We claim that  $f(x) = x^2$  is convex.

Pick  $a, b \in \mathfrak{R}$  and pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To check:

$$[\lambda a + (1 - \lambda)b]^2 \stackrel{?}{\leq} \lambda a^2 + (1 - \lambda)b^2.$$

We may assume that  $\lambda \neq 0, 1$ .

After simplifying

$$\lambda(1 - \lambda)2ab \stackrel{?}{\leq} \lambda(1 - \lambda)(a^2 + b^2),$$

## Example

We claim that  $f(x) = x^2$  is convex.

Pick  $a, b \in \mathfrak{R}$  and pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To check:

$$[\lambda a + (1 - \lambda)b]^2 \stackrel{?}{\leq} \lambda a^2 + (1 - \lambda)b^2.$$

We may assume that  $\lambda \neq 0, 1$ .

After simplifying

$$\lambda(1 - \lambda)2ab \stackrel{?}{\leq} \lambda(1 - \lambda)(a^2 + b^2),$$

or, equivalently, as  $\lambda, (1 - \lambda) > 0$ ,

$$a^2 + b^2 - 2ab \stackrel{?}{\geq} 0,$$

## Example

We claim that  $f(x) = x^2$  is convex.

Pick  $a, b \in \mathfrak{R}$  and pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To check:

$$[\lambda a + (1 - \lambda)b]^2 \stackrel{?}{\leq} \lambda a^2 + (1 - \lambda)b^2.$$

We may assume that  $\lambda \neq 0, 1$ .

After simplifying

$$\lambda(1 - \lambda)2ab \stackrel{?}{\leq} \lambda(1 - \lambda)(a^2 + b^2),$$

or, equivalently, as  $\lambda, (1 - \lambda) > 0$ ,

$$a^2 + b^2 - 2ab \stackrel{?}{\geq} 0,$$

which is the case as  $a^2 + b^2 - 2ab = (a - b)^2 \geq 0$ .

# Why Do We Care About Convex Functions?

## Proposition


Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .


Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.



## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .


Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.



## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.



## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

Let  $x = \lambda a + (1 - \lambda)b$ .

## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

Let  $x = \lambda a + (1 - \lambda)b$ .

Our goal is to show that  $x \in S$ , i.e., that  $g(x) \leq \beta$ .

## Proposition

Let  $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$  be a **convex** function and  $\beta \in \mathfrak{R}$ .

Then  $S = \{x \in \mathfrak{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

Let  $x = \lambda a + (1 - \lambda)b$ .

Our goal is to show that  $x \in S$ , i.e., that  $g(x) \leq \beta$ .

$$g(x) = g(\lambda a + (1 - \lambda)b)$$

## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

Let  $x = \lambda a + (1 - \lambda)b$ .

Our goal is to show that  $x \in S$ , i.e., that  $g(x) \leq \beta$ .

$$\begin{aligned} g(x) &= g(\lambda a + (1 - \lambda)b) \\ &\leq \lambda g(a) + (1 - \lambda)g(b) \quad (\text{convexity of } g) \end{aligned}$$

## Proposition

Let  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  be a **convex** function and  $\beta \in \mathbb{R}$ .

Then  $S = \{x \in \mathbb{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

Let  $x = \lambda a + (1 - \lambda)b$ .

Our goal is to show that  $x \in S$ , i.e., that  $g(x) \leq \beta$ .

$$\begin{aligned} g(x) &= g(\lambda a + (1 - \lambda)b) \\ &\leq \underbrace{\lambda}_{\geq 0} \underbrace{g(a)}_{\leq \beta} + \underbrace{(1 - \lambda)}_{\geq 0} \underbrace{g(b)}_{\leq \beta} \quad (\text{since } a, b \in S) \end{aligned}$$

## Proposition

Let  $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$  be a **convex** function and  $\beta \in \mathfrak{R}$ .

Then  $S = \{x \in \mathfrak{R}^n : g(x) \leq \beta\}$  is a **convex** set.

## Proof

Pick  $a, b \in S$ .

Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

Let  $x = \lambda a + (1 - \lambda)b$ .

Our goal is to show that  $x \in S$ , i.e., that  $g(x) \leq \beta$ .

$$\begin{aligned} g(x) &= g(\lambda a + (1 - \lambda)b) \\ &\leq \underbrace{\lambda}_{\geq 0} \underbrace{g(a)}_{\leq \beta} + \underbrace{(1 - \lambda)}_{\geq 0} \underbrace{g(b)}_{\leq \beta} && \text{(since } a, b \in S) \\ &\leq \lambda\beta + (1 - \lambda)\beta = \beta. \end{aligned}$$

## Proposition

$$\min c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

(P)

## Proposition

$$\min c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

(P)

If all functions  $g_i$  are **convex**, then the feasible region of (P) is **convex**.

## Proposition

$$\min c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

(P)

If all functions  $g_i$  are **convex**, then the feasible region of (P) is **convex**.

## Proof

## Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

If all functions  $g_i$  are **convex**, then the feasible region of (P) is **convex**.

## Proof

Let  $S_i = \{x : g_i(x) \leq 0\}$ .

## Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

If all functions  $g_i$  are **convex**, then the feasible region of (P) is **convex**.

## Proof

Let  $S_i = \{x : g_i(x) \leq 0\}$ .

By the previous result,  $S_i$  is convex.

## Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

(P)

If all functions  $g_i$  are **convex**, then the feasible region of (P) is **convex**.

## Proof

Let  $S_i = \{x : g_i(x) \leq 0\}$ .

By the previous result,  $S_i$  is convex.

The feasible region of (P) is  $S_1 \cap S_2 \cap \dots \cap S_k$ .

## Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array} \quad (\text{P})$$

If all functions  $g_i$  are **convex**, then the feasible region of (P) is **convex**.

## Proof

Let  $S_i = \{x : g_i(x) \leq 0\}$ .

By the previous result,  $S_i$  is convex.

The feasible region of (P) is  $S_1 \cap S_2 \cap \dots \cap S_k$ .

Since the intersection of convex sets is convex, the result follows.

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathcal{R}^n \rightarrow \mathcal{R}$  be a function.

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$  be a function. The **epigraph** of  $f$  is then given by,


$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathfrak{R}^n, y \in \mathfrak{R}, y \geq f(x) \right\} \subseteq \mathfrak{R}^{n+1}.$$

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$




# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$




$f$  is convex

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$



$f$  is convex

$\text{epi}(f)$  is convex

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$  be a function. The **epigraph** of  $f$  is then given by,


$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathfrak{R}^n, y \in \mathfrak{R}, y \geq f(x) \right\} \subseteq \mathfrak{R}^{n+1}.$$

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$




# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$




$f$  is NOT convex

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$



$f$  is NOT convex

$\text{epi}(f)$  is NOT convex

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

# Convex Functions Versus Convex Sets

## Definition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. The **epigraph** of  $f$  is then given by,

$$\text{epi}(f) = \left\{ \begin{pmatrix} y \\ x \end{pmatrix} : x \in \mathbb{R}^n, y \in \mathbb{R}, y \geq f(x) \right\} \subseteq \mathbb{R}^{n+1}.$$

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

1.  $f$  is convex  $\implies$   $\text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies$   $f$  is convex.

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix} \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ .

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix} \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix}, \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix}$$

## Proposition

Let  $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix}, \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} \lambda\alpha + (1 - \lambda)\beta \\ \lambda a + (1 - \lambda)b \end{pmatrix} \quad (\star)$$

## Proposition

Let  $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.
2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix}, \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} \lambda\alpha + (1 - \lambda)\beta \\ \lambda a + (1 - \lambda)b \end{pmatrix} \quad (\star)$$

Consider

$$f(\lambda a + (1 - \lambda)b)$$

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

1.  $f$  is convex  $\implies \text{epi}(f)$  is convex.

2.  $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix}, \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} \lambda\alpha + (1 - \lambda)\beta \\ \lambda a + (1 - \lambda)b \end{pmatrix} \quad (\star)$$

Consider

$$\begin{aligned} f(\lambda a + (1 - \lambda)b) &\leq && \text{(convexity of } f) \\ \lambda f(a) + (1 - \lambda)f(b) &&& \end{aligned}$$

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

- $f$  is convex  $\implies \text{epi}(f)$  is convex.
- $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix}, \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} \lambda\alpha + (1 - \lambda)\beta \\ \lambda a + (1 - \lambda)b \end{pmatrix} \quad (*)$$

Consider

$$\begin{aligned} f(\lambda a + (1 - \lambda)b) &\leq \quad (\text{convexity of } f) \\ \underbrace{\lambda}_{\geq 0} f(a) + \underbrace{(1 - \lambda)}_{\geq 0} f(b) \end{aligned}$$

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

- $f$  is convex  $\implies \text{epi}(f)$  is convex.
- $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix} \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} \lambda\alpha + (1 - \lambda)\beta \\ \lambda a + (1 - \lambda)b \end{pmatrix} \quad (*)$$

Consider

$$f(\lambda a + (1 - \lambda)b) \leq \underbrace{\lambda}_{\geq 0} \underbrace{f(a)}_{\leq \alpha} + \underbrace{(1 - \lambda)}_{\geq 0} \underbrace{f(b)}_{\leq \beta} \quad (\text{convexity of } f)$$

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

- $f$  is convex  $\implies \text{epi}(f)$  is convex.
- $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix}, \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} \lambda\alpha + (1 - \lambda)\beta \\ \lambda a + (1 - \lambda)b \end{pmatrix} \quad (*)$$

Consider

$$\begin{aligned} f(\lambda a + (1 - \lambda)b) &\leq \quad \text{(convexity of } f) \\ \underbrace{\lambda}_{\geq 0} \underbrace{f(a)}_{\leq \alpha} + \underbrace{(1 - \lambda)}_{\geq 0} \underbrace{f(b)}_{\leq \beta} &\leq \\ \lambda\alpha + (1 - \lambda)\beta. & \end{aligned}$$

## Proposition

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function. Then

- $f$  is convex  $\implies \text{epi}(f)$  is convex.
- $\text{epi}(f)$  is convex  $\implies f$  is convex.

## Proof

Pick  $\begin{pmatrix} \alpha \\ a \end{pmatrix}, \begin{pmatrix} \beta \\ b \end{pmatrix} \in \text{epi}(f)$ . Pick  $\lambda$  where  $0 \leq \lambda \leq 1$ .

To show:  $\text{epi}(f)$  contains

$$\lambda \begin{pmatrix} \alpha \\ a \end{pmatrix} + (1 - \lambda) \begin{pmatrix} \beta \\ b \end{pmatrix} = \begin{pmatrix} \lambda\alpha + (1 - \lambda)\beta \\ \lambda a + (1 - \lambda)b \end{pmatrix} \quad (*)$$

Consider

$$\begin{aligned} f(\lambda a + (1 - \lambda)b) &\leq \quad (\text{convexity of } f) \\ \underbrace{\lambda}_{\geq 0} \underbrace{f(a)}_{\leq \alpha} + \underbrace{(1 - \lambda)}_{\geq 0} \underbrace{f(b)}_{\leq \beta} &\leq \\ \lambda\alpha + (1 - \lambda)\beta. & \end{aligned}$$

Thus  $(*)$  is in  $\text{epi}(f)$ .

## Recap

## Recap

1. NLPs are hard in general.

## Recap

1. NLPs are hard in general.
2. We may assume the objective function of NLPs is linear.

## Recap

1. NLPs are hard in general.
2. We may assume the objective function of NLPs is linear.
3. Local optimum = optimal sol when the feasible region is convex.

## Recap

1. NLPs are hard in general.
2. We may assume the objective function of NLPs is linear.
3. Local optimum = optimal sol when the feasible region is convex.
4. We defined convex functions.

## Recap

1. NLPs are hard in general.
2. We may assume the objective function of NLPs is linear.
3. Local optimum = optimal sol when the feasible region is convex.
4. We defined convex functions.
5. Convex functions yield a convex feasible region.

## Recap

1. NLPs are hard in general.
2. We may assume the objective function of NLPs is linear.
3. Local optimum = optimal sol when the feasible region is convex.
4. We defined convex functions.
5. Convex functions yield a convex feasible region.
6. Convex functions and convex sets are related by epigraphs.