


## Module 3: Duality through examples (Weak Duality)

## Recap: Feasible Widths


Suppose we are given an instance of the **shortest path** problem...



## Recap: Feasible Widths

Suppose we are given an instance of the **shortest path** problem...


- a **graph**  $G = (V, E)$ ,
- a non-negative length  $c_e$  for each edge  $e \in E$ , and
- a pair of vertices  $s$  and  $t$  in  $V$ .



## Recap: Feasible Widths

Suppose we are given an instance of the **shortest path** problem...

- a **graph**  $G = (V, E)$ ,
- a non-negative length  $c_e$  for each edge  $e \in E$ , and
- a pair of vertices  $s$  and  $t$  in  $V$ .




A **width-assignment** is of the form

$$\{y_U : \delta(U) \text{ } s, t\text{-cut}\}$$

## Recap: Feasible Widths

Suppose we are given an instance of the **shortest path** problem...

- a **graph**  $G = (V, E)$ ,
- a non-negative length  $c_e$  for each edge  $e \in E$ , and
- a pair of vertices  $s$  and  $t$  in  $V$ .



A **width-assignment** is of the form


$$\{y_U : \delta(U) \text{ } s, t\text{-cut}\}$$

and this is **feasible** if the total width of cuts containing edge  $e$  is no more than  $c_e$ , for all  $e \in E$ .

## Recap: Feasible Widths

Suppose we are given an instance of the **shortest path** problem...

- a **graph**  $G = (V, E)$ ,
- a non-negative length  $c_e$  for each edge  $e \in E$ , and
- a pair of vertices  $s$  and  $t$  in  $V$ .



A **width-assignment** is of the form

$$\{y_U : \delta(U) \text{ } s, t\text{-cut}\}$$

and this is **feasible** if the total width of cuts containing edge  $e$  is no more than  $c_e$ , for all  $e \in E$ .

### Proposition

If  $y$  is a **feasible width assignment**, then any  $s, t$ -path must have length at least


$$\sum (y_U : U \text{ } s, t\text{-cut}).$$

## Recap: Feasible Widths

### Proposition

If  $y$  is a **feasible width assignment**, then any  $s, t$ -path must have length at least

$$\sum (y_U : U \text{ s,t-cut}).$$




Seemingly, we used an **ad hoc argument**, taylor-made for shortest paths...

## Recap: Feasible Widths

### Proposition

If  $y$  is a **feasible width assignment**, then any  $s, t$ -path must have length at least

$$\sum (y_U : U \text{ } s, t\text{-cut}).$$



Seemingly, we used an **ad hoc argument**, taylor-made for shortest paths. . .

but, as we will now see, there is a **constructive** and quite **mechanical** way to derive the Proposition via **linear programming**!

# An Instructive Example LP

The LP on the right is feasible...

$$\min (2, 3)x$$

$$\text{s.t. } \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

$$x \geq 0$$

# An Instructive Example LP

The LP on the right is feasible...

E.g.,  $x^1 = (8, 16)^\top$  and  
 $x^2 = (5, 13)^\top$  are feasible.

$$\begin{array}{ll} \min & (2, 3)x \\ \text{s.t.} & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{array}$$

# An Instructive Example LP

The LP on the right is feasible...

E.g.,  $x^1 = (8, 16)^\top$  and  
 $x^2 = (5, 13)^\top$  are feasible.

## Question

Can you find an **optimal** solution?

$$\begin{array}{ll} \min & (2, 3)x \\ \text{s.t.} & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{array}$$

# An Instructive Example LP

The LP on the right is feasible...

E.g.,  $x^1 = (8, 16)^\top$  and  
 $x^2 = (5, 13)^\top$  are feasible.

## Question

Can you find an **optimal** solution?

$x^1$  has an objective of value 64 and  
 $x^2$  has a value of 49  $\rightarrow$   $x^1$  is  
definitely not optimal, but is  $x^2$ ?

$$\begin{array}{ll} \min & (2, 3)x \\ \text{s.t.} & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{array}$$

# An Instructive Example LP

The LP on the right is feasible...

E.g.,  $x^1 = (8, 16)^\top$  and  
 $x^2 = (5, 13)^\top$  are feasible.

## Question

Can you find an **optimal** solution?

$x^1$  has an objective of value 64 and  
 $x^2$  has a value of 49  $\rightarrow x^1$  is  
definitely not optimal, but is  $x^2$ ?

$$\begin{array}{ll} \min & (2, 3)x \\ \text{s.t.} & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{array}$$

Feasible widths provide a  
**lower-bound** on the length of a  
shortest  $s, t$ -path...

# An Instructive Example LP

The LP on the right is feasible...

E.g.,  $x^1 = (8, 16)^\top$  and  
 $x^2 = (5, 13)^\top$  are feasible.

## Question

Can you find an **optimal** solution?

$x^1$  has an objective of value 64 and  
 $x^2$  has a value of 49  $\rightarrow x^1$  is  
definitely not optimal, but is  $x^2$ ?

$$\min (2, 3)x$$

$$\text{s.t.} \quad \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$
$$x \geq 0$$

Feasible widths provide a  
**lower-bound** on the length of a  
shortest  $s, t$ -path...

## Question

Can we find a good **lower-bound** on  
the objective value of the above LP?

## Deriving Valid Inequalities

Let's suppose that  $x$  is feasible for the LP on the right.

$$\begin{aligned} \min \quad & (2, 3)x \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

## Deriving Valid Inequalities

Let's suppose that  $x$  is feasible for the LP on the right.

It follows that  $x$  satisfies

$$\begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

$$\min (2, 3)x$$

$$\text{s.t. } \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$
$$x \geq 0$$

# Deriving Valid Inequalities

Let's suppose that  $x$  is feasible for the LP on the right.

It follows that  $x$  satisfies

$$\begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

and it also satisfies

$$\begin{array}{r} (2, 1)x \geq 20 \\ + (1, 1)x \geq 18 \\ + (-1, 1)x \geq 8 \\ \hline = (2, 3)x \geq 46 \end{array}$$

$$\min (2, 3)x$$

$$\text{s.t. } \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$
$$x \geq 0$$

# Deriving Valid Inequalities

Let's suppose that  $x$  is feasible for the LP on the right.

It follows that  $x$  satisfies

$$\begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

and it also satisfies

$$\begin{aligned} (2, 1)x &\geq 20 \\ + (1, 1)x &\geq 18 \\ + (-1, 1)x &\geq 8 \\ \hline &= (2, 3)x \geq 46 \end{aligned}$$

$$\min (2, 3)x$$

$$\begin{aligned} \text{s.t. } & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

Additionally, it satisfies

$$\begin{aligned} & y_1 \cdot (2, 1)x \geq y_1 \cdot 20 \\ & + y_2 \cdot (1, 1)x \geq y_2 \cdot 18 \\ & + y_3 \cdot (-1, 1)x \geq y_3 \cdot 8 \\ \hline & = (2y_1 + y_2 - y_3, y_1 + y_2 + y_3)x \\ & \geq 20y_1 + 18y_2 + 8y_3 \end{aligned}$$

for  $y_1, y_2, y_3 \geq 0$ .

So, if  $x$  is feasible for the LP on the right, it also satisfies

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

for any  $y_1, y_2, y_3 \geq 0$ .

$$\begin{aligned} \min \quad & (2, 3)x \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

So, if  $x$  is feasible for the LP on the right, it also satisfies

$$(y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

for any  $y_1, y_2, y_3 \geq 0$ .

E.g., for  $y = (0, 2, 1)^\top$ , we obtain

$$(1, 3)x \geq 44$$

$$\min (2, 3)x$$

$$\text{s.t. } \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$
$$x \geq 0$$

So, if  $x$  is feasible for the LP on the right, it also satisfies

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

for any  $y_1, y_2, y_3 \geq 0$ .

E.g., for  $y = (0, 2, 1)^\top$ , we obtain

$$(1, 3)x \geq 44$$

or

$$0 \geq 44 - (1, 3)x \quad (\star)$$

$$\min (2, 3)x$$

$$\text{s.t.} \quad \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$
$$x \geq 0$$

So, if  $x$  is feasible for the LP on the right, it also satisfies

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

for any  $y_1, y_2, y_3 \geq 0$ .

E.g., for  $y = (0, 2, 1)^\top$ , we obtain

$$(1, 3)x \geq 44$$

or

$$0 \geq 44 - (1, 3)x \quad (\star)$$

$$\min (2, 3)x$$

$$\text{s.t.} \quad \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$
$$x \geq 0$$

Therefore,

$$z(x) = (2, 3)x$$

So, if  $x$  is feasible for the LP on the right, it also satisfies

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

for any  $y_1, y_2, y_3 \geq 0$ .

E.g., for  $y = (0, 2, 1)^\top$ , we obtain

$$(1, 3)x \geq 44$$

or

$$0 \geq 44 - (1, 3)x \quad (\star)$$

$$\min (2, 3)x$$

$$\text{s.t.} \quad \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$
$$x \geq 0$$

Therefore,

$$\begin{aligned} z(x) &= (2, 3)x \\ &\geq (2, 3)x + 44 - (1, 3)x \\ &= 44 + (1, 0)x \end{aligned}$$

Since  $x \geq 0$ , it follows that

$$z(x) \geq 44$$

for every feasible solution  $x$ !

# State of Affairs

We now know that

$$\begin{aligned} \min \quad & (2, 3)x \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

# State of Affairs

We now know that

- (i)  $x^2 = (5, 13)^\top$  is a solution to the LP of **value 49** and

$$\begin{aligned} \min \quad & (2, 3)x \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

# State of Affairs

We now know that

- (i)  $x^2 = (5, 13)^\top$  is a solution to the LP of **value 49** and
- (ii)  $z(x) \geq 44$  for **every feasible solution** to the LP.

$$\begin{array}{ll} \min & (2, 3)x \\ \text{s.t.} & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{array}$$

# State of Affairs

We now know that

- (i)  $x^2 = (5, 13)^\top$  is a solution to the LP of **value 49** and
- (ii)  $z(x) \geq 44$  for **every feasible solution** to the LP.

→ The optimal value of the LP is in the interval  $[44, 49]$ .

$$\begin{array}{ll} \min & (2, 3)x \\ \text{s.t.} & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{array}$$

# State of Affairs

We now know that

- (i)  $x^2 = (5, 13)^\top$  is a solution to the LP of **value 49** and
- (ii)  $z(x) \geq 44$  for **every feasible solution** to the LP.

→ The optimal value of the LP is in the interval  $[44, 49]$ .

Can we find a better **lowerbound** on  $z(x)$  for a feasible  $x$ ?

$$\begin{array}{ll} \min & (2, 3)x \\ \text{s.t.} & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{array}$$

# Lowerbounding $z(x)$ **Systematically!**

We know that a feasible  $x$  satisfies

$$0 \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} - (y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x$$

for any  $y_1, y_2, y_3 \geq 0$ .

## Lowerbounding $z(x)$ **Systematically!**

We know that a feasible  $x$  satisfies

$$0 \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} - (y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x$$

for any  $y_1, y_2, y_3 \geq 0$ . **Therefore,**

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} + \left( (2, 3) - (y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \right) x \quad (\star)$$

# Lowerbounding $z(x)$ **Systematically!**

We know that a feasible  $x$  satisfies

$$0 \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} - (y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x$$

for any  $y_1, y_2, y_3 \geq 0$ . **Therefore,**

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} + \left( (2, 3) - (y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \right) x \quad (\star)$$

We want the second term to be **non-negative**. Since  $x \geq 0$ , this amounts to choosing  $y$  such that

$$(y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \leq (2, 3)$$

# Lowerbounding $z(x)$ **Systematically!**

We know that a feasible  $x$  satisfies

$$0 \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} - (y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x$$

for any  $y_1, y_2, y_3 \geq 0$ . **Therefore,**

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} + \left( (2, 3) - (y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \right) x \quad (\star)$$

We want the second term to be **non-negative**. Since  $x \geq 0$ , this amounts to choosing  $y$  such that

$$(y_1, y_2, y_2) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \leq (2, 3)$$

With such a  $y$  we then have from  $(\star)$ :

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix}$$

## Lowerbounding $z(x)$ **Systematically!**

So, we choose  $y \geq 0$  such that

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \leq (2, 3) \quad (\star)$$

yields

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \quad (\diamond)$$

# Lowerbounding $z(x)$ **Systematically!**

So, we choose  $y \geq 0$  such that

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \leq (2, 3) \quad (\star)$$

yields

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \quad (\diamond)$$

## Idea

Find the best possible lower-bound on  $z$ .

# Lowerbounding $z(x)$ **Systematically!**

So, we choose  $y \geq 0$  such that

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \leq (2, 3) \quad (\star)$$

yields

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \quad (\diamond)$$

## Idea

Find the best possible lower-bound on  $z$ . I.e., find  $y \geq 0$  such that  $(\star)$  holds, and the right-hand side of  $(\diamond)$  is maximized!

# Lowerbounding $z(x)$ **Systematically!**

So, we choose  $y \geq 0$  such that

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} \leq (2, 3) \quad (\star)$$

yields

$$z(x) \geq (y_1, y_2, y_3) \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \quad (\diamond)$$

## Idea

Find the best possible lower-bound on  $z$ . I.e., find  $y \geq 0$  such that  $(\star)$  holds, and the right-hand side of  $(\diamond)$  is maximized!

This is a **Linear Program**:

$$\begin{aligned} \max \quad & (20, 18, 8)y \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} y \leq (2, 3) \\ & y \geq 0 \end{aligned}$$

# Lowerbounding $z(x)$ Systematically!

This is a **Linear Program**:

$$\begin{aligned} \max \quad & (20, 18, 8)y \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} y \leq (2, 3) \\ & y \geq 0 \end{aligned}$$

Solving it gives:

$$\begin{aligned} \bar{y}_1 &= 0 \\ \bar{y}_2 &= 5/2 \\ \bar{y}_3 &= 1/2 \end{aligned}$$

and the **objective value** is 49.

# Lowerbounding $z(x)$ **Systematically!**

This is a **Linear Program**:

$$\begin{aligned} \max \quad & (20, 18, 8)y \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} y \leq (2, 3) \\ & y \geq 0 \end{aligned}$$

**Solving it gives:**

$$\begin{aligned} \bar{y}_1 &= 0 \\ \bar{y}_2 &= 5/2 \\ \bar{y}_3 &= 1/2 \end{aligned}$$

and the **objective value** is 49.

There is **no feasible solution**  $x$  to

$$\begin{aligned} \min \quad & (2, 3)x \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

which has an objective value smaller than 49.

# Lowerbounding $z(x)$ **Systematically!**

This is a **Linear Program**:

$$\begin{aligned} \max \quad & (20, 18, 8)y \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} y \leq (2, 3) \\ & y \geq 0 \end{aligned}$$

**Solving it gives:**

$$\begin{aligned} \bar{y}_1 &= 0 \\ \bar{y}_2 &= 5/2 \\ \bar{y}_3 &= 1/2 \end{aligned}$$

and the **objective value** is 49.

There is **no feasible solution**  $x$  to

$$\begin{aligned} \min \quad & (2, 3)x \\ \text{s.t.} \quad & \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \geq \begin{pmatrix} 20 \\ 18 \\ 8 \end{pmatrix} \\ & x \geq 0 \end{aligned}$$

which has an objective value smaller than 49.

Since  $x^2 = (5, 13)^\top$  is a feasible solution with value 49, it **must be optimal!**

# A General Argument

Suppose now we are given the LP

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & Ax \geq b \\ & x \geq 0 \end{array}$$

# A General Argument

Suppose now we are given the LP

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & Ax \geq b \\ & x \geq 0 \end{array}$$

Any **feasible** solution  $x$  must satisfy

$$y^\top Ax \geq y^\top b,$$

for  $y \geq 0$ ,

# A General Argument

Suppose now we are given the LP

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & Ax \geq b \\ & x \geq 0 \end{array}$$

Any **feasible** solution  $x$  must satisfy

$$y^\top Ax \geq y^\top b,$$

for  $y \geq 0$ , and hence also

$$0 \geq y^\top b - y^\top Ax$$

# A General Argument

Suppose now we are given the LP

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & Ax \geq b \\ & x \geq 0 \end{array}$$

Therefore,

$$\begin{aligned} z(x) &= c^\top x \\ &\geq c^\top x + y^\top b - y^\top Ax \end{aligned}$$

Any **feasible** solution  $x$  must satisfy

$$y^\top Ax \geq y^\top b,$$

for  $y \geq 0$ , and hence also

$$0 \geq y^\top b - y^\top Ax$$

# A General Argument

Suppose now we are given the LP

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & Ax \geq b \\ & x \geq 0 \end{array}$$

Any **feasible** solution  $x$  must satisfy

$$y^\top Ax \geq y^\top b,$$

for  $y \geq 0$ , and hence also

$$0 \geq y^\top b - y^\top Ax$$

Therefore,

$$\begin{aligned} z(x) &= c^\top x \\ &\geq c^\top x + y^\top b - y^\top Ax \\ &= y^\top b + (c^\top - y^\top A)x \end{aligned}$$

# A General Argument

Suppose now we are given the LP

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & Ax \geq b \\ & x \geq 0 \end{array}$$

Any **feasible** solution  $x$  must satisfy

$$y^\top Ax \geq y^\top b,$$

for  $y \geq 0$ , and hence also

$$0 \geq y^\top b - y^\top Ax$$

Therefore,

$$\begin{aligned} z(x) &= c^\top x \\ &\geq c^\top x + y^\top b - y^\top Ax \\ &= y^\top b + (c^\top - y^\top A)x \end{aligned}$$

If we also know that

$$A^\top y \leq c$$

then  $x \geq 0$  implies that  $z(x) \geq y^\top b$ .

# A General Argument

Suppose now we are given the LP

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & Ax \geq b \\ & x \geq 0 \end{array}$$

Any **feasible** solution  $x$  must satisfy

$$y^\top Ax \geq y^\top b,$$

for  $y \geq 0$ , and hence also

$$0 \geq y^\top b - y^\top Ax$$

Therefore,

$$\begin{aligned} z(x) &= c^\top x \\ &\geq c^\top x + y^\top b - y^\top Ax \\ &= y^\top b + (c^\top - y^\top A)x \end{aligned}$$

If we also know that

$$A^\top y \leq c$$

then  $x \geq 0$  implies that  $z(x) \geq y^\top b$ .

The best **lower-bound on  $z(x)$**  can be found by the following LP:

$$\begin{array}{ll} \max & b^\top y \\ \text{s.t.} & A^\top y \leq c \\ & y \geq 0 \end{array}$$

# The Dual LP

The linear program

$$\max \quad b^T y \quad (D)$$

$$\text{s.t.} \quad A^T y \leq c$$

$$y \geq 0$$

is called the **dual** of **primal** LP

$$\min \quad c^T x \quad (P)$$

$$\text{s.t.} \quad Ax \geq b$$

$$x \geq 0$$

# The Dual LP

The linear program

$$\begin{aligned} \max \quad & b^T y && \text{(D)} \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

is called the **dual** of **primal** LP

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq b \\ & x \geq 0 \end{aligned}$$

## Theorem

[**Weak Duality**] If  $\bar{x}$  is feasible for (P) and  $\bar{y}$  is feasible for (D), then  $b^T \bar{y} \leq c^T \bar{x}$ .

# The Dual LP

The linear program

$$\begin{aligned} \max \quad & b^T y & (D) \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

is called the **dual** of **primal** LP

$$\begin{aligned} \min \quad & c^T x & (P) \\ \text{s.t.} \quad & Ax \geq b \\ & x \geq 0 \end{aligned}$$

## Theorem

[**Weak Duality**] If  $\bar{x}$  is feasible for (P) and  $\bar{y}$  is feasible for (D), then  $b^T \bar{y} \leq c^T \bar{x}$ .

**Proof:**

$$b^T \bar{y} = \bar{y}^T b$$

# The Dual LP

The linear program

$$\begin{aligned} \max \quad & b^T y & (D) \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

is called the **dual** of **primal** LP

$$\begin{aligned} \min \quad & c^T x & (P) \\ \text{s.t.} \quad & Ax \geq b \\ & x \geq 0 \end{aligned}$$

## Theorem

[**Weak Duality**] If  $\bar{x}$  is feasible for (P) and  $\bar{y}$  is feasible for (D), then  $b^T \bar{y} \leq c^T \bar{x}$ .

**Proof:**

$$b^T \bar{y} = \bar{y}^T b \leq \bar{y}^T (A\bar{x}) = (A^T \bar{y})^T \bar{x}$$

as  $\bar{y} \geq 0$  and  $b \leq A\bar{x}$ ,

# The Dual LP

The linear program

$$\begin{aligned} \max \quad & b^T y & (D) \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

is called the **dual** of **primal** LP

$$\begin{aligned} \min \quad & c^T x & (P) \\ \text{s.t.} \quad & Ax \geq b \\ & x \geq 0 \end{aligned}$$

## Theorem

[**Weak Duality**] If  $\bar{x}$  is feasible for (P) and  $\bar{y}$  is feasible for (D), then  $b^T \bar{y} \leq c^T \bar{x}$ .

**Proof:**

$$b^T \bar{y} = \bar{y}^T b \leq \bar{y}^T (A\bar{x}) = (A^T \bar{y})^T \bar{x} \leq c^T \bar{x}$$

as  $\bar{y} \geq 0$  and  $b \leq A\bar{x}$ , as  $\bar{x} \geq 0$  and  $A^T \bar{y} \leq c$ . □

Lowerbounding the Length of  $s, t$ -Paths

## Recap: Shortest Path LP

Given a **shortest path** instance  $G = (V, E)$ ,  $s, t \in V$ ,  $c_e \geq 0$  for all  $e \in E$ , the shortest-path LP is

$$\min \sum (c_e x_e : e \in E)$$

$$\text{s.t. } \sum (x_e : e \in \delta(U)) \geq 1 \quad (U \subseteq V, s \in U, t \notin U)$$

$$x \geq 0, x \text{ integer}$$

## Recap: Shortest Path LP


Given a **shortest path** instance  $G = (V, E)$ ,  $s, t \in V$ ,  $c_e \geq 0$  for all  $e \in E$ , the shortest-path LP is

$$\begin{aligned} \min \quad & \sum (c_e x_e : e \in E) \\ \text{s.t.} \quad & \sum (x_e : e \in \delta(U)) \geq 1 \quad (U \subseteq V, s \in U, t \notin U) \\ & x \geq 0, x \text{ integer} \end{aligned}$$

Let's look at an **example**!

## Shortest Path: **Example**

On the right, we see a sample instance of the shortest-path problem.



## Shortest Path: Example


On the right, we see a sample instance of the shortest-path problem.

Here is the corresponding IP:

$$\min (3, 4, 1, 2, 2)x$$

$$\text{s.t.} \quad \begin{array}{l} \{s\} \\ \{s, a\} \\ \{s, b\} \\ \{s, a, b\} \end{array} \begin{pmatrix} sa & sb & ab & at & bt \\ \left( \begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right) x \geq \mathbb{1} \end{pmatrix}$$

$$x \geq 0, x \text{ integer}$$



## Shortest Path: Example


On the right, we see a sample instance of the shortest-path problem.

Here is the corresponding IP:

$$\min (3, 4, 1, 2, 2)x$$

$$\text{s.t.} \quad \begin{array}{l} \{s\} \\ \{s, a\} \\ \{s, b\} \\ \{s, a, b\} \end{array} \begin{pmatrix} sa & sb & ab & at & bt \\ \left( \begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right) \end{pmatrix} x \geq \mathbb{1}$$

$$x \geq 0, x \text{ integer}$$



Note that if  $P$  is an  $s, t$ -path, then letting

$$\bar{x}_e = \begin{cases} 1 & \text{if } e \text{ is an edge of } P \\ 0 & \text{otherwise.} \end{cases}$$

for all  $e \in E$  yields a feasible IP solution and

## Shortest Path: Example


On the right, we see a sample instance of the shortest-path problem.

Here is the corresponding IP:

$$\min (3, 4, 1, 2, 2)x$$

$$\text{s.t.} \quad \begin{array}{l} \{s\} \\ \{s, a\} \\ \{s, b\} \\ \{s, a, b\} \end{array} \begin{pmatrix} sa & sb & ab & at & bt \\ \left( \begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right) \end{pmatrix} x \geq \mathbb{1}$$

$$x \geq 0, x \text{ integer}$$



**Note** that if  $P$  is an  $s, t$ -path, then letting


$$\bar{x}_e = \begin{cases} 1 & \text{if } e \text{ is an edge of } P \\ 0 & \text{otherwise.} \end{cases}$$

for all  $e \in E$  yields a **feasible** IP solution and its **objective value** is  $c(P)$ .

$$\min (3, 4, 1, 2, 2)x$$

$$\text{s.t.} \quad \begin{matrix} & sa & sb & ab & at & bt \\ \{s\} & \left( \begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right) & x & \geq & \mathbb{1} \\ \{s, a\} & & & & & \\ \{s, b\} & & & & & \\ \{s, a, b\} & & & & & \end{matrix}$$

$$x \geq 0, x \text{ integer}$$



Example:


$$P = sa, ab, bt$$

is an  $s, t$ -path.

$$\min (3, 4, 1, 2, 2)x$$

$$\text{s.t.} \quad \begin{array}{l} \{s\} \\ \{s, a\} \\ \{s, b\} \\ \{s, a, b\} \end{array} \begin{pmatrix} sa & sb & ab & at & bt \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{pmatrix} x \geq \mathbb{1}$$

$$x \geq 0, x \text{ integer}$$



**Example:**

$$P = sa, ab, bt$$

is an  $s, t$ -path.


$$x = (1, 0, 1, 0, 1)^T$$

is feasible for the IP, and its value is 6.

$$\min (3, 4, 1, 2, 2)x$$

$$\text{s.t.} \quad \begin{matrix} & sa & sb & ab & at & bt \\ \{s\} & \left( \begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right) & x & \geq & \mathbb{1} \\ \{s, a\} & & & & & \\ \{s, b\} & & & & & \\ \{s, a, b\} & & & & & \end{matrix}$$

$$x \geq 0, x \text{ integer}$$



**Example:**

$$P = sa, ab, bt$$

is an  $s, t$ -path.

$$x = (1, 0, 1, 0, 1)^T$$

is feasible for the IP, and its value is 6.


## Remark

The optimal value of the shortest path IP is, at most, the length of a shortest  $s, t$ -path.

$$\min (3, 4, 1, 2, 2)x \quad (\text{P})$$

$$\text{s.t.} \quad \begin{matrix} & sa & sb & ab & at & bt \\ \{s\} & \left( \begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right) & x \geq \mathbb{1} \\ \{s, a\} \\ \{s, b\} \\ \{s, a, b\} \end{matrix}$$

$$x \geq 0, x \text{ integer}$$




**Note** that dropping the **integrality** restriction can not increase the optimal value.

min  $(3, 4, 1, 2, 2)x$  (P)

$$\text{s.t.} \quad \begin{array}{l} \{s\} \\ \{s, a\} \\ \{s, b\} \\ \{s, a, b\} \end{array} \begin{pmatrix} sa & sb & ab & at & bt \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{pmatrix} x \geq \mathbb{1}$$

$x \geq 0, x$  integer




**Note** that dropping the **integrality** restriction can not increase the optimal value.

The resulting LP is called the **linear programming relaxation** of the IP.

$$\min (3, 4, 1, 2, 2)x \quad (\text{P})$$

$$\text{s.t.} \quad \begin{matrix} & sa & sb & ab & at & bt \\ \{s\} & \left( \begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right) & x & \geq & \mathbb{1} \\ \{s, a\} & & & & & \\ \{s, b\} & & & & & \\ \{s, a, b\} & & & & & \end{matrix}$$

$$x \geq 0, x \text{ integer}$$



**Note** that dropping the **integrality** restriction can not increase the optimal value.

The resulting LP is called the **linear programming relaxation** of the IP.


Straight from **Weak Duality** theorem, we have that:

### Remark

The dual of (P) has optimal value no larger than that of (P)!


The **dual** of the shortest path LP on the previous slide is given by

$$\begin{array}{l}
 \max \quad \mathbb{1}^\top y \\
 \text{s.t.} \quad \begin{array}{cccc}
 & \{s\} & \{s, a\} & \{s, b\} & \{s, a, b\} \\
 sa & \left( \begin{array}{cccc}
 1 & 0 & 1 & 0 \\
 1 & 1 & 0 & 0 \\
 0 & 1 & 1 & 0 \\
 0 & 1 & 0 & 1 \\
 0 & 0 & 1 & 1
 \end{array} \right) & y \leq & \begin{pmatrix} 3 \\ 4 \\ 1 \\ 2 \\ 2 \end{pmatrix} \\
 ab & & & & \\
 at & & & & \\
 bt & & & & \\
 y \geq & \mathbb{0} & & & 
 \end{array}
 \end{array}$$



The **dual** of the shortest path LP on the previous slide is given by


$$\begin{array}{l}
 \max \quad \mathbb{1}^\top y \\
 \text{s.t.} \quad \begin{array}{cccc}
 & \{s\} & \{s, a\} & \{s, b\} & \{s, a, b\} \\
 sa & \left( \begin{array}{cccc}
 1 & 0 & 1 & 0 \\
 1 & 1 & 0 & 0 \\
 0 & 1 & 1 & 0 \\
 0 & 1 & 0 & 1 \\
 0 & 0 & 1 & 1
 \end{array} \right) & y \leq & \begin{pmatrix} 3 \\ 4 \\ 1 \\ 2 \\ 2 \end{pmatrix} \\
 bt & & & & \\
 y \geq & \mathbb{0} & & & 
 \end{array}
 \end{array}$$



**Note** that dual solutions assign the value  $y_U \geq 0$  to every  $s, t$ -cut  $\delta(U)$ !

The **dual** of the shortest path LP on the previous slide is given by

$$\begin{array}{l}
 \max \quad \mathbb{1}^\top y \\
 \text{s.t.} \quad \begin{array}{cccc}
 & \{s\} & \{s, a\} & \{s, b\} & \{s, a, b\} \\
 sa & \left( \begin{array}{cccc}
 1 & 0 & 1 & 0 \\
 1 & 1 & 0 & 0 \\
 0 & 1 & 1 & 0 \\
 0 & 1 & 0 & 1 \\
 0 & 0 & 1 & 1
 \end{array} \right) & y \leq & \begin{pmatrix} 3 \\ 4 \\ 1 \\ 2 \\ 2 \end{pmatrix} \\
 y \geq 0
 \end{array}
 \end{array}$$




**Note** that dual solutions assign the value  $y_U \geq 0$  to every  $s, t$ -cut  $\delta(U)$ !

Focus on the constraint for **edge**  $ab$ :

$$y_{\{s,a\}} + y_{\{s,b\}} \leq 1$$

The **dual** of the shortest path LP on the previous slide is given by

$$\begin{array}{l} \max \quad \mathbb{1}^\top y \\ \\ \text{s.t.} \quad \begin{array}{cccc} & \{s\} & \{s, a\} & \{s, b\} & \{s, a, b\} \\ sa & \left( \begin{array}{cccc} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{array} \right) & y \leq & \begin{pmatrix} 3 \\ 4 \\ 1 \\ 2 \\ 2 \end{pmatrix} \\ y \geq \mathbb{0} \end{array} \end{array}$$



**Note** that dual solutions assign the value  $y_U \geq 0$  to every  $s, t$ -cut  $\delta(U)$ !


Focus on the constraint for **edge  $ab$** :

$$y_{\{s,a\}} + y_{\{s,b\}} \leq 1$$

The left-hand side is precisely the  $y$ -value assigned to  $s, t$ -cuts containing  $ab$ !

The **dual** of the shortest path LP on the previous slide is given by

$$\begin{array}{l}
 \max \quad \mathbb{1}^\top y \\
 \\
 \text{s.t.} \quad \begin{array}{cccc}
 & \{s\} & \{s, a\} & \{s, b\} & \{s, a, b\} \\
 sa & \left( \begin{array}{cccc}
 1 & 0 & 1 & 0 \\
 1 & 1 & 0 & 0 \\
 0 & 1 & 1 & 0 \\
 0 & 1 & 0 & 1 \\
 0 & 0 & 1 & 1
 \end{array} \right) & y \leq & \begin{pmatrix} 3 \\ 4 \\ 1 \\ 2 \\ 2 \end{pmatrix} \\
 \\
 y \geq 0
 \end{array}
 \end{array}$$



**Note** that dual solutions assign the value  $y_U \geq 0$  to every  $s, t$ -cut  $\delta(U)$ !

## Remark

$y$  is feasible for the above LP if and only if it is a feasible width assignment for the  $s, t$ -cuts in the given shortest path instance!

# General Shortest Path Instances

**Input:**  $G = (V, E)$ ,  $c_e \geq 0$  for all  $e \in E$ ,  $s, t \in V$ .

# General Shortest Path Instances

**Input:**  $G = (V, E)$ ,  $c_e \geq 0$  for all  $e \in E$ ,  $s, t \in V$ .

Shortest path LP:

$$\begin{aligned} \min \quad & \sum (c_e x_e : e \in E) \\ \text{s.t.} \quad & \sum (x_e : e \in \delta(U)) \geq 1 \\ & (\delta(U) \text{ } s, t \text{ - cut}) \\ & x \geq 0 \end{aligned}$$

# General Shortest Path Instances

**Input:**  $G = (V, E)$ ,  $c_e \geq 0$  for all  $e \in E$ ,  $s, t \in V$ .

Shortest path LP:

$$\begin{aligned} \min \quad & \sum (c_e x_e : e \in E) \\ \text{s.t.} \quad & \sum (x_e : e \in \delta(U)) \geq 1 \\ & (\delta(U) \text{ } s, t \text{ - cut}) \\ & x \geq 0 \end{aligned}$$

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbf{1} \\ & x \geq 0 \end{aligned}$$

where

# General Shortest Path Instances

**Input:**  $G = (V, E)$ ,  $c_e \geq 0$  for all  $e \in E$ ,  $s, t \in V$ .

Shortest path LP:

$$\begin{aligned} \min \quad & \sum (c_e x_e : e \in E) \\ \text{s.t.} \quad & \sum (x_e : e \in \delta(U)) \geq 1 \\ & (\delta(U) \text{ } s, t \text{-cut}) \\ & x \geq 0 \end{aligned}$$

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbf{1} \\ & x \geq 0 \end{aligned}$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .

# General Shortest Path Instances

**Input:**  $G = (V, E)$ ,  $c_e \geq 0$  for all  $e \in E$ ,  $s, t \in V$ .

Shortest path LP:

$$\begin{aligned} \min \quad & \sum (c_e x_e : e \in E) \\ \text{s.t.} \quad & \sum (x_e : e \in \delta(U)) \geq 1 \\ & (\delta(U) \text{ } s, t \text{-cut}) \\ & x \geq 0 \end{aligned}$$

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbf{1} \\ & x \geq 0 \end{aligned}$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .
- (ii)  $A[U, e] = 1$  if  $e \in \delta(U)$  and 0 otherwise.

The LP is of the form

$$\min \quad c^T x \quad (\text{P})$$

$$\text{s.t.} \quad Ax \geq \mathbb{1} \\ x \geq 0$$

Its dual is of the form

$$\max \quad \mathbb{1}^T y \quad (\text{D})$$

$$\text{s.t.} \quad A^T y \leq c \\ y \geq 0$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .
- (ii)  $A[U, e] = 1$  if  $e \in \delta(U)$ , and 0 otherwise.

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbb{1} \\ & x \geq 0 \end{aligned}$$

Its dual is of the form

$$\begin{aligned} \max \quad & \mathbb{1}^T y && \text{(D)} \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .
- (ii)  $A[U, e] = 1$  if  $e \in \delta(U)$ , and 0 otherwise.

**Note** that the dual has a constraint for every edge  $e \in E$ .

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbf{1} \\ & x \geq 0 \end{aligned}$$

Its dual is of the form

$$\begin{aligned} \max \quad & \mathbf{1}^T y && \text{(D)} \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .
- (ii)  $A[U, e] = 1$  if  $e \in \delta(U)$ , and 0 otherwise.

**Note** that the dual has a constraint for every edge  $e \in E$ . The left-hand side of this constraint is

$$\sum (y_U : e \in \delta(U))$$

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbf{1} \\ & x \geq 0 \end{aligned}$$

Its dual is of the form

$$\begin{aligned} \max \quad & \mathbf{1}^T y && \text{(D)} \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .
- (ii)  $A[U, e] = 1$  if  $e \in \delta(U)$ , and 0 otherwise.

**Note** that the dual has a constraint for every edge  $e \in E$ . The left-hand side of this constraint is

$$\sum (y_U : e \in \delta(U))$$

and the right-hand side is  $c_e$ .

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbf{1} \\ & x \geq 0 \end{aligned}$$

Its dual is of the form

$$\begin{aligned} \max \quad & \mathbf{1}^T y && \text{(D)} \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .
- (ii)  $A[U, e] = 1$  if  $e \in \delta(U)$ , and 0 otherwise.

**Note** that the dual has a constraint for every edge  $e \in E$ . The left-hand side of this constraint is

$$\sum (y_U : e \in \delta(U))$$

and the right-hand side is  $c_e$ .

## Remark

Feasible solutions to (D) correspond precisely to feasible width assignments.

The LP is of the form

$$\begin{aligned} \min \quad & c^T x && \text{(P)} \\ \text{s.t.} \quad & Ax \geq \mathbf{1} \\ & x \geq 0 \end{aligned}$$

Its dual is of the form

$$\begin{aligned} \max \quad & \mathbf{1}^T y && \text{(D)} \\ \text{s.t.} \quad & A^T y \leq c \\ & y \geq 0 \end{aligned}$$

where

- (i)  $A$  has a column for every edge and a row for every  $s, t$ -cut  $\delta(U)$ .
- (ii)  $A[U, e] = 1$  if  $e \in \delta(U)$ , and 0 otherwise.

**Note** that the dual has a constraint for every edge  $e \in E$ . The left-hand side of this constraint is

$$\sum (y_U : e \in \delta(U))$$

and the right-hand side is  $c_e$ .

## Remark

Feasible solutions to (D) correspond precisely to feasible width assignments. **Weak duality** implies that  $\sum y_U$  is, at most, the length of a shortest  $s, t$ -path!

## Recap

- The **dual LP** of

$$\min\{c^T x : Ax \geq b, x \geq 0\} \quad (\text{P})$$

is given by

$$\max\{b^T y : A^T y \leq c, y \geq 0\} \quad (\text{D})$$

## Recap

- The **dual LP** of

$$\min\{c^T x : Ax \geq b, x \geq 0\} \quad (\text{P})$$

is given by

$$\max\{b^T y : A^T y \leq c, y \geq 0\} \quad (\text{D})$$

- If  $x$  is feasible for (P) and  $y$  feasible for (D), then  $b^T y \leq c^T x$ .

## Recap

- The **dual LP** of

$$\min\{c^T x : Ax \geq b, x \geq 0\} \quad (\text{P})$$

is given by

$$\max\{b^T y : A^T y \leq c, y \geq 0\} \quad (\text{D})$$

- If  $x$  is feasible for (P) and  $y$  feasible for (D), then  $b^T y \leq c^T x$ .
- The **LP relaxation** of an integer program is obtained by dropping the **integrality restriction**.

## Recap

- The **dual LP** of

$$\min\{c^T x : Ax \geq b, x \geq 0\} \quad (\text{P})$$

is given by

$$\max\{b^T y : A^T y \leq c, y \geq 0\} \quad (\text{D})$$

- If  $x$  is feasible for (P) and  $y$  feasible for (D), then  $b^T y \leq c^T x$ .
- The **LP relaxation** of an integer program is obtained by dropping the **integrality restriction**.
- The **dual** of the **shortest path LP** is given by

$$\begin{aligned} \max \quad & \sum (y_U : \delta(U) \text{ s, t-cut}) \\ \text{s.t.} \quad & \sum (y_U : e \in \delta(U)) \leq c_e \quad (e \in E) \\ & y \geq 0 \end{aligned}$$