

Module 6: Nonlinear Programs (the KKT theorem)

$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



$$(1) \quad x_2 \geq x_1^2;$$


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



$$(1) \quad x_2 \geq x_1^2;$$

$$(2) \quad x_1 \geq x_2^2;$$


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



$$(1) \quad x_2 \geq x_1^2;$$

$$(2) \quad x_1 \geq x_2^2;$$

$$(3) \quad x_1 \geq \frac{1}{2}.$$


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$




$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to the NLP.


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to the NLP.

How do we prove this?


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to the NLP.

How do we prove this?

Step 1. Find a relaxation of the NLP.


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to the NLP.

How do we prove this?

Step 1. Find a relaxation of the NLP.

Step 2. Prove \bar{x} is optimal for the relaxation.


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to the NLP.

How do we prove this?

Step 1. Find a relaxation of the NLP.

Step 2. Prove \bar{x} is optimal for the relaxation.

Step 3. Deduce that \bar{x} is optimal for the NLP.

Original NLP


$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$



Original NLP


$$\min -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$




Relaxation

$$\min -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$



Original NLP


$$\min -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$




New relaxation

$$\min -x_1 - x_2$$

s.t.


$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$




Original NLP

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$



New relaxation

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & 2x_1 - x_2 \leq 1 \quad (a) \\ & -x_1 + 2x_2 \leq 1 \quad (b) \end{array}$$



Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to

$$\min \quad -x_1 - x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to

$$\max x_1 + x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to

$$\max x_1 + x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Proof

Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to

$$\max x_1 + x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Proof

Tight constraints for \bar{x} are (a) and (b).

Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to

$$\max x_1 + x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Proof

Tight constraints for \bar{x} are (a) and (b).

Goal: Show that the objective function is in the cone of tight constraints.

Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to

$$\max x_1 + x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Proof

Tight constraints for \bar{x} are (a) and (b).

Goal: Show that the objective function is in the cone of tight constraints.

$$\begin{pmatrix} 1 \\ 1 \end{pmatrix} \stackrel{?}{\in} \text{cone} \left\{ \begin{pmatrix} 2 \\ -1 \end{pmatrix}, \begin{pmatrix} -1 \\ 2 \end{pmatrix} \right\}$$

Claim

$\bar{x} = (1, 1)^\top$ is an optimal solution to

$$\max x_1 + x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Proof

Tight constraints for \bar{x} are (a) and (b).

Goal: Show that the objective function is in the cone of tight constraints.

$$\begin{pmatrix} 1 \\ 1 \end{pmatrix} \stackrel{?}{\in} \text{cone} \left\{ \begin{pmatrix} 2 \\ -1 \end{pmatrix}, \begin{pmatrix} -1 \\ 2 \end{pmatrix} \right\} \quad \Leftarrow$$

$$\begin{pmatrix} 1 \\ 1 \end{pmatrix} = 1 \times \begin{pmatrix} 2 \\ -1 \end{pmatrix} + 1 \times \begin{pmatrix} -1 \\ 2 \end{pmatrix} \quad \checkmark$$

Original NLP

$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$

Relaxation

$$\min \quad -x_1 - x_2$$

s.t.

$$2x_1 - x_2 \leq 1 \quad (a)$$

$$-x_1 + 2x_2 \leq 1 \quad (b)$$

Original NLP

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

Relaxation

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & 2x_1 - x_2 \leq 1 \quad (a) \\ & -x_1 + 2x_2 \leq 1 \quad (b) \end{array}$$

$\bar{x} = (1, 1)^\top$ is an optimal solution to the relaxation

Original NLP

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

Relaxation

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & 2x_1 - x_2 \leq 1 \quad (a) \\ & -x_1 + 2x_2 \leq 1 \quad (b) \end{array}$$

$\bar{x} = (1, 1)^\top$ is an optimal solution to the relaxation

 \bar{x} is an optimal solution to the *original NLP*

Original NLP

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

Relaxation

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & 2x_1 - x_2 \leq 1 \quad (a) \\ & -x_1 + 2x_2 \leq 1 \quad (b) \end{array}$$

$\bar{x} = (1, 1)^\top$ is an optimal solution to the relaxation



\bar{x} is an optimal solution to the *original NLP*

Question

Can we do this in general?

Original NLP

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

Relaxation

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & 2x_1 - x_2 \leq 1 \quad (a) \\ & -x_1 + 2x_2 \leq 1 \quad (b) \end{array}$$

$\bar{x} = (1, 1)^\top$ is an optimal solution to the relaxation



\bar{x} is an optimal solution to the *original NLP*

Question

Can we do this in general? YES

Original NLP

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

Relaxation

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & 2x_1 - x_2 \leq 1 \quad (a) \\ & -x_1 + 2x_2 \leq 1 \quad (b) \end{array}$$

$\bar{x} = (1, 1)^\top$ is an optimal solution to the relaxation



\bar{x} is an optimal solution to the *original NLP*

Question

Can we do this in general? **YES**

The key tool we'll use is **subgradients**.

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

$$h(\bar{x}) = f(\bar{x})$$

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

$$h(\bar{x}) = f(\bar{x})$$

h is a lower bound for f

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.


Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

$$h(\bar{x}) = f(\bar{x})$$

h is a lower bound for f



Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if


$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

$$h(\bar{x}) = f(\bar{x})$$

h is a lower bound for f

unique subgradient



Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if


$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

$$h(\bar{x}) = f(\bar{x})$$

h is a lower bound for f

first subgradient



Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if


$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

$$h(\bar{x}) = f(\bar{x})$$

h is a lower bound for f

second subgradient



Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and $\bar{x} \in \mathbb{R}^n$.

Then, $s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if


$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

$h(x)$ is affine

$$h(\bar{x}) = f(\bar{x})$$

h is a lower bound for f

NOT UNIQUE



Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$h(x) = f(\bar{x}) + s^\top(x - \bar{x}) =$$

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$\begin{aligned} h(x) &= f(\bar{x}) + s^\top(x - \bar{x}) = \\ &= 0 + \end{aligned}$$

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$\begin{aligned} h(x) &= f(\bar{x}) + s^\top (x - \bar{x}) = \\ &= 0 + (-1, 2)(x - (1, 1)^\top) = \end{aligned}$$

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$\begin{aligned} h(x) &= f(\bar{x}) + s^\top(x - \bar{x}) = \\ &= 0 + (-1, 2)(x - (1, 1)^\top) = -x_1 + 2x_2 - 1. \end{aligned}$$

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$\begin{aligned} h(x) &= f(\bar{x}) + s^\top (x - \bar{x}) = \\ &= 0 + (-1, 2)(x - (1, 1)^\top) = -x_1 + 2x_2 - 1. \end{aligned}$$

Check: $h(x) \leq f(x)$ for all $x \in \mathbb{R}^n$.

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top(x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$\begin{aligned} h(x) &= f(\bar{x}) + s^\top(x - \bar{x}) = \\ &= 0 + (-1, 2)(x - (1, 1)^\top) = -x_1 + 2x_2 - 1. \end{aligned}$$

Check: $h(x) \leq f(x)$ for all $x \in \mathbb{R}^n$.

$$-x_1 + 2x_2 - 1 \stackrel{?}{\leq} -x_1 + x_2^2$$

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$\begin{aligned} h(x) &= f(\bar{x}) + s^\top (x - \bar{x}) = \\ &= 0 + (-1, 2)(x - (1, 1)^\top) = -x_1 + 2x_2 - 1. \end{aligned}$$

Check: $h(x) \leq f(x)$ for all $x \in \mathbb{R}^n$.

$$-x_1 + 2x_2 - 1 \stackrel{?}{\leq} -x_1 + x_2^2$$

or equivalently,

$$x_2^2 - 2x_2 + 1 \stackrel{?}{\geq} 0,$$

Definition

$s \in \mathbb{R}^n$ is a **subgradient** of f at \bar{x} if

$$h(x) := f(\bar{x}) + s^\top (x - \bar{x}) \leq f(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Example

Consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ where $f(x) = -x_1 + x_2^2$ and $\bar{x} = (1, 1)^\top$.

We claim that $(-1, 2)^\top$ is a subgradient of f at \bar{x} .

$$\begin{aligned} h(x) &= f(\bar{x}) + s^\top (x - \bar{x}) = \\ &= 0 + (-1, 2)(x - (1, 1)^\top) = -x_1 + 2x_2 - 1. \end{aligned}$$

Check: $h(x) \leq f(x)$ for all $x \in \mathbb{R}^n$.

$$-x_1 + 2x_2 - 1 \stackrel{?}{\leq} -x_1 + x_2^2$$

or equivalently,

$$x_2^2 - 2x_2 + 1 \stackrel{?}{\geq} 0,$$

which is the case as $x_2^2 - 2x_2 + 1 = (x_2 - 1)^2 \geq 0$.

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if


- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .




Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .




UNIQUE SUPPORTING HALFSPACE AT \bar{x} .

Definition

Let $C \in \mathbb{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .




Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .




Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .



NON-UNIQUE SUPPORTING HALFSPACE AT \bar{x} .

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .

Definition

Let $C \in \mathbb{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .

Question

What do we get when $n = 1$?

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .

Question

What do we get when $n = 1$?

- C is a segment (or a halfline)

Definition

Let $C \in \mathfrak{R}^n$ be a convex set and let $\bar{x} \in C$.

The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .

Question

What do we get when $n = 1$?

- C is a segment (or a halfline)
- F is a halfline

Definition

Let $C \in \mathbb{R}^n$ be a convex set and let $\bar{x} \in C$.


The halfspace $F = \{x : s^\top x \leq \beta\}$ is **supporting** C at \bar{x} if

- (1) $C \subseteq F$ and
- (2) $s^\top \bar{x} = \beta$. That is, \bar{x} is on the boundary of F .

Question

What do we get when $n = 1$?

- C is a segment (or a halfline)
- F is a halfline



Subgradients and Supporting Halfspaces

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{K}^n \rightarrow \mathfrak{K}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Remark

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Remark

- C is convex, as g is a convex function,

Subgradients and Supporting Halfspaces

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Remark

- C is convex, as g is a convex function,
- F is a halfspace, as $h(x)$ is an affine function, and

Subgradients and Supporting Halfspaces

Proposition

Let $g : \Re^n \rightarrow \Re$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Remark

- C is convex, as g is a convex function,
- F is a halfspace, as $h(x)$ is an affine function, and
- $h(\bar{x}) = g(\bar{x}) = 0$ thus, \bar{x} is on the boundary of F .

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Example

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Example

$$g(x) = x_2^2 - x_1$$

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .


Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Example

$$g(x) = x_2^2 - x_1$$



Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.


Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Example

$$g(x) = x_2^2 - x_1$$

$$\bar{x} = (1, 1)^\top$$



Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.


Then, F is a supporting halfspace of C at \bar{x} .

Example

$$g(x) = x_2^2 - x_1$$

$$\bar{x} = (1, 1)^\top$$

$$s = (-1, 2)^\top \text{ subgradient at } \bar{x}$$



Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .


Example

$$g(x) = x_2^2 - x_1$$

$$\bar{x} = (1, 1)^\top$$

$$s = (-1, 2)^\top \text{ subgradient at } \bar{x}$$

$$\begin{aligned} h(x) &= 0 + (-1, 2) \left[\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right] \\ &= -x_1 + 2x_2 - 1 \end{aligned}$$



Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Example


$$g(x) = x_2^2 - x_1$$

$$\bar{x} = (1, 1)^\top$$

$$s = (-1, 2)^\top \text{ subgradient at } \bar{x}$$

$$\begin{aligned} h(x) &= 0 + (-1, 2) \left[\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right] \\ &= -x_1 + 2x_2 - 1 \end{aligned}$$

$$F = \{x : -x_1 + 2x_2 \leq 1\}$$



Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition


Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .



Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Let $x \in C$ and thus, $g(x) \leq 0$

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Let $x \in C$ and thus, $g(x) \leq 0$

By definition of a subgradient, we know that $h(x) \leq g(x)$.

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Let $x \in C$ and thus, $g(x) \leq 0$

By definition of a subgradient, we know that $h(x) \leq g(x)$.

It follows that $h(x) \leq g(x) \leq 0$.

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Let $x \in C$ and thus, $g(x) \leq 0$

By definition of a subgradient, we know that $h(x) \leq g(x)$.

It follows that $h(x) \leq g(x) \leq 0$.

Hence, $x \in F$.

Proposition

Let $g : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Let $x \in C$ and thus, $g(x) \leq 0$

By definition of a subgradient, we know that $h(x) \leq g(x)$.

It follows that $h(x) \leq g(x) \leq 0$.

Hence, $x \in F$.

Claim: $h(\bar{x}) = 0$

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Let $x \in C$ and thus, $g(x) \leq 0$

By definition of a subgradient, we know that $h(x) \leq g(x)$.

It follows that $h(x) \leq g(x) \leq 0$.

Hence, $x \in F$.

Claim: $h(\bar{x}) = 0$

$h(\bar{x}) = g(\bar{x})$

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Proof

Claim: $C \subseteq F$.

Let $x \in C$ and thus, $g(x) \leq 0$

By definition of a subgradient, we know that $h(x) \leq g(x)$.

It follows that $h(x) \leq g(x) \leq 0$.

Hence, $x \in F$.

Claim: $h(\bar{x}) = 0$

$h(\bar{x}) = g(\bar{x}) = 0$.

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Question

Why is this relevant for us?

Proposition

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and let \bar{x} where $g(\bar{x}) = 0$.

Let s be a subgradient of g at \bar{x} .

Let $C = \{x : g(x) \leq 0\}$.

Let $F = \{x : h(x) := g(\bar{x}) + s^\top(x - \bar{x}) \leq 0\}$.

Then, F is a supporting halfspace of C at \bar{x} .

Question

Why is this relevant for us?



WE USE IT TO CONSTRUCT RELAXATIONS OF NLPs

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

\bar{x} is a feasible solution

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

\bar{x} is a feasible solution

g_1 is convex

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

\bar{x} is a feasible solution

g_1 is convex

$$g_1(\bar{x}) = 0$$

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

\bar{x} is a feasible solution

g_1 is convex

$$g_1(\bar{x}) = 0$$

s is a subgradient for g_1 at \bar{x}

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

\bar{x} is a feasible solution

g_1 is convex

$$g_1(\bar{x}) = 0$$

s is a subgradient for g_1 at \bar{x}

If we replace the nonlinear constraint

$$g_1(x) \leq 0$$

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

\bar{x} is a feasible solution

g_1 is convex

$$g_1(\bar{x}) = 0$$

s is a subgradient for g_1 at \bar{x}

If we replace the nonlinear constraint

$$g_1(x) \leq 0$$

with the linear constraint

$$h(x) = g_1(\bar{x}) + s^\top(x - \bar{x}) \leq 0$$

we get a relaxation.

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

\bar{x} is a feasible solution

g_1 is convex

$$g_1(\bar{x}) = 0$$

s is a subgradient for g_1 at \bar{x}


If we replace the nonlinear constraint

$$g_1(x) \leq 0$$

with the linear constraint

$$h(x) = g_1(\bar{x}) + s^\top(x - \bar{x}) \leq 0$$

we get a relaxation.



Proposition

$$\min c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Example

$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$

$\bar{x} = (1, 1)^\top$ feasible

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Example

$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$

$\bar{x} = (1, 1)^\top$ feasible

$$I = \{1, 2\}$$

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Example

$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$

$\bar{x} = (1, 1)^\top$ feasible

$$I = \{1, 2\}$$

$(2, -1)^\top$ subgradient for g_1 at \bar{x}

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Example

$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$

$\bar{x} = (1, 1)^\top$ feasible

$$I = \{1, 2\}$$

$(2, -1)^\top$ subgradient for g_1 at \bar{x}

$(-1, 2)^\top$ subgradient for g_2 at \bar{x}

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Example

$$\min \quad -x_1 - x_2$$

s.t.

$$-x_2 + x_1^2 \leq 0 \quad (1)$$

$$-x_1 + x_2^2 \leq 0 \quad (2)$$

$$-x_1 + \frac{1}{2} \leq 0 \quad (3)$$

$\bar{x} = (1, 1)^\top$ feasible

$$I = \{1, 2\}$$

$(2, -1)^\top$ subgradient for g_1 at \bar{x}

$(-1, 2)^\top$ subgradient for g_2 at \bar{x}

$$-\begin{pmatrix} -1 \\ -1 \end{pmatrix} \in \text{cone} \left\{ \begin{pmatrix} 2 \\ -1 \end{pmatrix}, \begin{pmatrix} -1 \\ 2 \end{pmatrix} \right\} \quad \Rightarrow \quad \bar{x} \text{ optimal.}$$

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i \in I)$$

Proposition

$$\min c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\min c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i \in I)$$

We proved that the set of solutions to $g_i(x) \leq 0$

is contained in the set of solutions to $g_i(\bar{x}) + s^{(i)}(x - \bar{x}) \leq 0$.

Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(\bar{x}) + s^{(i)}(x - \bar{x}) \leq 0 \quad (i \in I) \end{array}$$

Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(\bar{x}) + s^{(i)}(x - \bar{x}) \leq 0 \quad (i \in I) \end{array}$$

$g_i(\bar{x}) + s^{(i)}(x - \bar{x}) \leq 0$ can be rewritten as

Proposition

$$\min c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\min c^\top x$$

s.t.

$$g_i(\bar{x}) + s^{(i)}(x - \bar{x}) \leq 0 \quad (i \in I)$$

$g_i(\bar{x}) + s^{(i)}(x - \bar{x}) \leq 0$ can be rewritten as

$$s^{(i)}x \leq s^{(i)}\bar{x} - g_i(\bar{x})$$

Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\begin{array}{ll} \max & -c^\top x \\ \text{s.t.} & \\ & s^{(i)}x \leq s^{(i)}\bar{x} - g_i(\bar{x}) \quad (i \in I) \end{array}$$

Proposition

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\max \quad -c^\top x$$

s.t.

$$s^{(i)}x \leq s^{(i)}\bar{x} - g_i(\bar{x}) \quad (i \in I)$$

Then, \bar{x} is optimal for the relaxation if $-c \in \text{cone} \{s^{(i)} : i \in I\}$.

Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\begin{array}{ll} \max & -c^\top x \\ \text{s.t.} & \\ & s^{(i)}x \leq s^{(i)}\bar{x} - g_i(\bar{x}) \quad (i \in I) \end{array}$$

Then, \bar{x} is optimal for the relaxation if $-c \in \text{cone} \{s^{(i)} : i \in I\}$.

This means that \bar{x} is also optimal for the NLP.

Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\begin{array}{ll} \max & -c^\top x \\ \text{s.t.} & \\ & s^{(i)}x \leq s^{(i)}\bar{x} - g_i(\bar{x}) \quad (i \in I) \end{array}$$

Then, \bar{x} is optimal for the relaxation if $-c \in \text{cone} \{s^{(i)} : i \in I\}$.

This means that \bar{x} is also optimal for the NLP.

Question

Is there a converse to this result?

Proposition

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

g_1, \dots, g_k all convex

\bar{x} is a feasible solution

$$\forall i \in I, g_i(\bar{x}) = 0$$

$\forall i \in I, s^{(i)}$ subgradient for g_i at \bar{x}

If $-c \in \text{cone} \{s^{(i)} : i \in I\}$ then \bar{x} is **optimal**.

Proof

We have a relaxation

$$\begin{array}{ll} \max & -c^\top x \\ \text{s.t.} & \\ & s^{(i)}x \leq s^{(i)}\bar{x} - g_i(\bar{x}) \quad (i \in I) \end{array}$$

Then, \bar{x} is optimal for the relaxation if $-c \in \text{cone} \{s^{(i)} : i \in I\}$.

This means that \bar{x} is also optimal for the NLP.

Question

Is there a converse to this result? **YES**

Gradients: A Calculus Detour

Gradients: A Calculus Detour

Proposition

Let $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be a convex function and let $\bar{x} \in \mathfrak{R}^n$.

Gradients: A Calculus Detour

Proposition

Let $f : \mathfrak{R}^n \rightarrow \mathfrak{R}$ be a convex function and let $\bar{x} \in \mathfrak{R}^n$.

If the **gradient** $\nabla f(\bar{x})$ of f exists at \bar{x} , then it is a subgradient.

Gradients: A Calculus Detour

Proposition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function and let $\bar{x} \in \mathbb{R}^n$.

If the **gradient** $\nabla f(\bar{x})$ of f exists at \bar{x} , then it is a subgradient.

Proposition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be function and let $\bar{x} \in \mathbb{R}^n$.

If the partial derivative $\frac{\partial f(x)}{\partial x_j}$ exists for f at \bar{x} for all $j = 1, \dots, n$, then the gradient $\nabla f(\bar{x})$ is obtained by evaluating for \bar{x} ,

$$\left(\frac{\partial f(x)}{\partial x_1}, \dots, \frac{\partial f(x)}{\partial x_n} \right)^\top.$$

Example

Compute the gradient of the convex function

$$f(x) = -x_2 + x_1^2$$

at $\bar{x} = (1, 1)^\top$.

Example

Compute the gradient of the convex function

$$f(x) = -x_2 + x_1^2$$

at $\bar{x} = (1, 1)^\top$.

We have

$$\left(\frac{\partial f(x)}{\partial x_1}, \frac{\partial f(x)}{\partial x_2} \right)^\top = (2x_1, -1)^\top$$

Example

Compute the gradient of the convex function

$$f(x) = -x_2 + x_1^2$$

at $\bar{x} = (1, 1)^\top$.

We have

$$\left(\frac{\partial f(x)}{\partial x_1}, \frac{\partial f(x)}{\partial x_2} \right)^\top = (2x_1, -1)^\top$$

For \bar{x} we get $\nabla f(\bar{x}) = (2, -1)^\top$.

Example

Compute the gradient of the convex function

$$f(x) = -x_2 + x_1^2$$

at $\bar{x} = (1, 1)^\top$.

We have

$$\left(\frac{\partial f(x)}{\partial x_1}, \frac{\partial f(x)}{\partial x_2} \right)^\top = (2x_1, -1)^\top$$

For \bar{x} we get $\nabla f(\bar{x}) = (2, -1)^\top$.

Since $(2, -1)^\top$ is the gradient of f at \bar{x} , it is a subgradient as well.

Definition

A feasible solution to \bar{x} is a **Slater point** of

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

if $g_i(\bar{x}) < 0$ for all $i = 1, \dots, k$.

Definition

A feasible solution to \bar{x} is a **Slater point** of

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

if $g_i(\bar{x}) < 0$ for all $i = 1, \dots, k$.

Example

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

Definition

A feasible solution to \bar{x} is a **Slater point** of

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

if $g_i(\bar{x}) < 0$ for all $i = 1, \dots, k$.

Example

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

$\bar{x} = \left(\frac{3}{4}, \frac{3}{4}\right)^\top$ is a Slater point.

Definition

A feasible solution to \bar{x} is a **Slater point** of


$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

if $g_i(\bar{x}) < 0$ for all $i = 1, \dots, k$.

Example

$$\begin{array}{ll} \min & -x_1 - x_2 \\ \text{s.t.} & \\ & -x_2 + x_1^2 \leq 0 \quad (1) \\ & -x_1 + x_2^2 \leq 0 \quad (2) \\ & -x_1 + \frac{1}{2} \leq 0 \quad (3) \end{array}$$

$\bar{x} = \left(\frac{3}{4}, \frac{3}{4}\right)^\top$ is a Slater point.



The Karush-Kuhn-Tucker (KKT) Theorem

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\min \quad c^\top x$$

s.t.

$$g_i(x) \leq 0 \quad (i = 1, \dots, k)$$

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

1. g_1, \dots, g_k are all convex,

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

1. g_1, \dots, g_k are all convex,
2. there exists a **Slater point**,

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

1. g_1, \dots, g_k are all convex,
2. there exists a **Slater point**,
3. \bar{x} is a feasible solution,

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

1. g_1, \dots, g_k are all convex,
2. there exists a **Slater point**,
3. \bar{x} is a feasible solution,
4. I is the set of indices i for which $g_i(\bar{x}) = 0$, and

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

1. g_1, \dots, g_k are all convex,
2. there exists a **Slater point**,
3. \bar{x} is a feasible solution,
4. I is the set of indices i for which $g_i(\bar{x}) = 0$, and
5. for all $i \in I$ there exists a gradient $\nabla g_i(\bar{x})$ of g_i at \bar{x} .

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

1. g_1, \dots, g_k are all convex,
2. there exists a **Slater point**,
3. \bar{x} is a feasible solution,
4. I is the set of indices i for which $g_i(\bar{x}) = 0$, and
5. for all $i \in I$ there exists a gradient $\nabla g_i(\bar{x})$ of g_i at \bar{x} .

Then \bar{x} is optimal $\iff -c \in \text{cone} \{ \nabla g_i(\bar{x}) : i \in I \}$.

The Karush-Kuhn-Tucker (KKT) Theorem

Consider the following NLP:

$$\begin{array}{ll} \min & c^\top x \\ \text{s.t.} & \\ & g_i(x) \leq 0 \quad (i = 1, \dots, k) \end{array}$$

Suppose that

1. g_1, \dots, g_k are all convex,
2. there exists a **Slater point**,
3. \bar{x} is a feasible solution,
4. I is the set of indices i for which $g_i(\bar{x}) = 0$, and
5. for all $i \in I$ there exists a gradient $\nabla g_i(\bar{x})$ of g_i at \bar{x} .

Then \bar{x} is optimal $\iff -c \in \text{cone} \{ \nabla g_i(\bar{x}) : i \in I \}$.

Remark

We proved the “easy” direction “ \iff ”.

Recap

Recap

- We showed how to prove optimality using relaxations.

Recap

- We showed how to prove optimality using relaxations.
- We defined subgradients.

Recap

- We showed how to prove optimality using relaxations.
- We defined subgradients.
- We defined supporting halfspaces.

Recap

- We showed how to prove optimality using relaxations.
- We defined subgradients.
- We defined supporting halfspaces.
- We related subgradients and supporting halfspaces.

Recap

- We showed how to prove optimality using relaxations.
- We defined subgradients.
- We defined supporting halfspaces.
- We related subgradients and supporting halfspaces.
- We showed how to relax convex constraints by a linear constraint.

Recap

- We showed how to prove optimality using relaxations.
- We defined subgradients.
- We defined supporting halfspaces.
- We related subgradients and supporting halfspaces.
- We showed how to relax convex constraints by a linear constraint.
- We gave sufficient conditions for a solution to be optimal.

Recap

- We showed how to prove optimality using relaxations.
- We defined subgradients.
- We defined supporting halfspaces.
- We related subgradients and supporting halfspaces.
- We showed how to relax convex constraints by a linear constraint.
- We gave sufficient conditions for a solution to be optimal.
- We stated the KKT theorem.