

Module 2: Linear Programs (Canonical Forms)

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Definition

Let B be a basis of A . Then (P) is in **canonical form** for B if

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Definition

Let B be a basis of A . Then (P) is in **canonical form** for B if

(P1) $A_B = I$, and

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Definition

Let B be a basis of A . Then (P) is in **canonical form** for B if

(P1) $A_B = I$, and

(P2) $c_j = 0$ for all $j \in B$.

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Definition

Let B be a basis of A . Then (P) is in **canonical form** for B if

(P1) $A_B = I$, and

(P2) $c_j = 0$ for all $j \in B$.

$$\max \quad (0 \quad 0 \quad 2 \quad 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

Canonical form for
 $B = \{1, 2\}$

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Definition

Let B be a basis of A . Then (P) is in **canonical form** for B if

(P1) $A_B = I$, and

(P2) $c_j = 0$ for all $j \in B$.

$$\max \quad (-2 \quad 0 \quad 0 \quad 6)x + 2$$

s.t.

$$\begin{pmatrix} -1 & 1 & 0 & 3 \\ 1 & 0 & 1 & -1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

Canonical form for
 $B = \{2, 3\}$

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Idea

For any basis B we can “rewrite” (P) so that it is in canonical form for a basis B and such that the resulting LP behaves the same as (P).

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Idea

For any basis B we can “rewrite” (P) so that it is in canonical form for a basis B and such that the resulting LP behaves the same as (P).

More formally, we will show the following:

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Idea

For any basis B we can “rewrite” (P) so that it is in canonical form for a basis B and such that the resulting LP behaves the same as (P).

More formally, we will show the following:

Proposition

For any basis B , there exists (P') in canonical form for B such that

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Idea

For any basis B we can “rewrite” (P) so that it is in canonical form for a basis B and such that the resulting LP behaves the same as (P).

More formally, we will show the following:

Proposition

For any basis B , there exists (P') in canonical form for B such that
(1) (P) and (P') have the same feasible region, and

Consider

$$\max \{c^\top x : Ax = b, x \geq 0\} \quad (\text{P})$$

Idea

For any basis B we can “rewrite” (P) so that it is in canonical form for a basis B and such that the resulting LP behaves the same as (P).

More formally, we will show the following:

Proposition

For any basis B , there exists (P') in canonical form for B such that

- (1) (P) and (P') have the same feasible region, and
- (2) feasible solutions have the same objective value for (P) and (P').

Proposition

For any basis B , there exists (P') in canonical form for B such that

- (1) (P) and (P') have the same feasible region, and
- (2) feasible solutions have the same objective value for (P) and (P') .

Proposition

For any basis B , there exists (P') in canonical form for B such that

(1) (P) and (P') have the same feasible region, and

(2) feasible solutions have the same objective value for (P) and (P') .

$$\max \quad (0 \quad 0 \quad 2 \quad 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

Proposition

For any basis B , there exists (P') in canonical form for B such that

- (1) (P) and (P') have the same feasible region, and
- (2) feasible solutions have the same objective value for (P) and (P') .

$$\max \quad (0 \quad 0 \quad 2 \quad 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$\max \quad (-2 \quad 0 \quad 0 \quad 6)x + 2$$

s.t.

$$\begin{pmatrix} -1 & 1 & 0 & 3 \\ 1 & 0 & 1 & -1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

Proposition

For any basis B , there exists (P') in canonical form for B such that

- (1) (P) and (P') have the same feasible region, and
- (2) feasible solutions have the same objective value for (P) and (P') .

$\max \quad (0 \quad 0 \quad 2 \quad 4)x$	$\max \quad (-2 \quad 0 \quad 0 \quad 6)x + 2$
s.t.	s.t.
$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$	$\begin{pmatrix} -1 & 1 & 0 & 3 \\ 1 & 0 & 1 & -1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$
$x_1, x_2, x_3, x_4 \geq 0$	$x_1, x_2, x_3, x_4 \geq 0$

Example

- (1) $\bar{x} = (1, 2, 0, 0)^\top$ is feasible for both LPs.

Proposition

For any basis B , there exists (P') in canonical form for B such that

- (1) (P) and (P') have the same feasible region, and
- (2) feasible solutions have the same objective value for (P) and (P') .

$\begin{aligned} \max \quad & (0 \quad 0 \quad 2 \quad 4)x \\ \text{s.t.} \quad & \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ & x_1, x_2, x_3, x_4 \geq 0 \end{aligned}$	$\begin{aligned} \max \quad & (-2 \quad 0 \quad 0 \quad 6)x + 2 \\ \text{s.t.} \quad & \begin{pmatrix} -1 & 1 & 0 & 3 \\ 1 & 0 & 1 & -1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \\ & x_1, x_2, x_3, x_4 \geq 0 \end{aligned}$
--	--

Example

- (1) $\bar{x} = (1, 2, 0, 0)^\top$ is feasible for both LPs.

- (2)
$$\begin{aligned} (0 \quad 0 \quad 2 \quad 4)\bar{x} &= 2 \times 0 + 4 \times 0 = 0 \\ (-2 \quad 0 \quad 0 \quad 6)\bar{x} + 2 &= -2 \times 1 + 6 \times 0 + 2 = 0 \end{aligned}$$

Illustration with an Example

Illustration with an Example

$$\begin{array}{ll} \max & \underbrace{(0 \quad 0 \quad 2 \quad 4)}_c x \\ \text{s.t.} & \\ & \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ & x_1, x_2, x_3, x_4 \geq 0 \end{array}$$

(P)

Illustration with an Example

$$\begin{array}{ll} \max & \underbrace{(0 \quad 0 \quad 2 \quad 4)}_c x \\ \text{s.t.} & \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ & x_1, x_2, x_3, x_4 \geq 0 \end{array} \quad (\text{P})$$

Question

How do we rewrite (P) in canonical form for basis $B = \{2, 3\}$?

Illustration with an Example

$$\begin{array}{ll} \max & \underbrace{(0 \quad 0 \quad 2 \quad 4)}_c x \\ \text{s.t.} & \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ & x_1, x_2, x_3, x_4 \geq 0 \end{array} \quad (\text{P})$$

Question

How do we rewrite (P) in canonical form for basis $B = \{2, 3\}$?

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_{\{2,3\}} = I$.

Illustration with an Example

$$\begin{array}{ll} \max & \underbrace{(0 \quad 0 \quad 2 \quad 4)}_c x \\ \text{s.t.} & \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ & x_1, x_2, x_3, x_4 \geq 0 \end{array} \quad (\text{P})$$

Question

How do we rewrite (P) in canonical form for basis $B = \{2, 3\}$?

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_{\{2,3\}} = I$.

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ where $\bar{c}_2 = \bar{c}_3 = 0$ and \bar{z} is a constant.

Rewriting Constraints – Example

Rewriting Constraints – Example

$$\underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b$$

Rewriting Constraints – Example

$$\underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b$$

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_{\{2,3\}} = I$.

Rewriting Constraints – Example

$$\underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b$$

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_{\{2,3\}} = I$.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Rewriting Constraints – Example

$$\underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b$$

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_{\{2,3\}} = I$.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$



$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Rewriting Constraints – Example

$$\underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b$$

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_{\{2,3\}} = I$.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$



$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$



$$\begin{pmatrix} -1 & 1 & 0 & 3 \\ 1 & 0 & 1 & -1 \end{pmatrix} x = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Rewriting Constraints – General

Rewriting Constraints – General

Consider the system $Ax = b$ with basis B of A .

Rewriting Constraints – General

Consider the system $Ax = b$ with basis B of A .

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_B = I$ for some basis B .

Rewriting Constraints – General

Consider the system $Ax = b$ with basis B of A .

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_B = I$ for some basis B .

$$Ax = b$$

Rewriting Constraints – General

Consider the system $Ax = b$ with basis B of A .

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_B = I$ for some basis B .

$$Ax = b$$



$$\underbrace{A_B^{-1}A}_{A'}x = \underbrace{A_B^{-1}b}_{b'}$$

Rewriting Constraints – General

Consider the system $Ax = b$ with basis B of A .

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_B = I$ for some basis B .

$$Ax = b$$



$$\underbrace{A_B^{-1}A}_{A'}x = \underbrace{A_B^{-1}b}_{b'}$$

Remarks

- $A'_B = I$.

Rewriting Constraints – General

Consider the system $Ax = b$ with basis B of A .

(P1) Replace $Ax = b$ by $A'x = b'$ with $A'_B = I$ for some basis B .

$$Ax = b$$



$$\underbrace{A_B^{-1}A}_{A'}x = \underbrace{A_B^{-1}b}_{b'}$$

Remarks

- $A'_B = I$.
- $Ax = b$ and $A'x = b'$ have the same set of solutions.

Rewriting the Objective Function – Example

Rewriting the Objective Function – Example

$$\max \quad z = \underbrace{(0 \quad 0 \quad 2 \quad 4)}_{c^T} x$$

s.t.

$$\underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b$$

$$x_1, x_2, x_3, x_4 \geq 0$$

Rewriting the Objective Function – Example

$$\begin{array}{l} \max \quad z = \underbrace{(0 \quad 0 \quad 2 \quad 4)}_{c^\top} x \\ \text{s.t.} \\ \quad \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ \quad x_1, x_2, x_3, x_4 \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ where $\bar{c}_2 = \bar{c}_3 = 0$ and \bar{z} is a constant.

Rewriting the Objective Function – Example

$$\begin{array}{l} \max \quad z = \underbrace{(0 \quad 0 \quad 2 \quad 4)}_{c^\top} x \\ \text{s.t.} \\ \quad \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ \quad x_1, x_2, x_3, x_4 \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ where $\bar{c}_2 = \bar{c}_3 = 0$ and \bar{z} is a constant.

Step 1. Construct a new objective function by

Rewriting the Objective Function – Example

$$\begin{array}{l} \max \quad z = \underbrace{(0 \quad 0 \quad 2 \quad 4)}_{c^\top} x \\ \text{s.t.} \\ \quad \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ \quad x_1, x_2, x_3, x_4 \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ where $\bar{c}_2 = \bar{c}_3 = 0$ and \bar{z} is a constant.

Step 1. Construct a new objective function by

- multiplying constraint 1 by y_1 ,

Rewriting the Objective Function – Example

$$\begin{array}{l} \max \quad z = \underbrace{(0 \quad 0 \quad 2 \quad 4)}_{c^\top} x \\ \text{s.t.} \\ \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ x_1, x_2, x_3, x_4 \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ where $\bar{c}_2 = \bar{c}_3 = 0$ and \bar{z} is a constant.

Step 1. Construct a new objective function by

- multiplying constraint 1 by y_1 ,
- multiplying constraint 2 by y_2 , and

Rewriting the Objective Function – Example

$$\begin{array}{l} \max \quad z = \underbrace{(0 \quad 0 \quad 2 \quad 4)}_{c^\top} x \\ \text{s.t.} \\ \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ x_1, x_2, x_3, x_4 \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ where $\bar{c}_2 = \bar{c}_3 = 0$ and \bar{z} is a constant.

Step 1. Construct a new objective function by

- multiplying constraint 1 by y_1 ,
- multiplying constraint 2 by y_2 , and
- adding the resulting constraints to the objective function.

Rewriting the Objective Function – Example

$$\begin{array}{l} \max \quad z = \underbrace{(0 \quad 0 \quad 2 \quad 4)}_{c^\top} x \\ \text{s.t.} \\ \underbrace{\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix}}_A x = \underbrace{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}_b \\ x_1, x_2, x_3, x_4 \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ where $\bar{c}_2 = \bar{c}_3 = 0$ and \bar{z} is a constant.

Step 1. Construct a new objective function by

- multiplying constraint 1 by y_1 ,
- multiplying constraint 2 by y_2 , and
- adding the resulting constraints to the objective function.

Step 2. Choose y_1, y_2 to get $\bar{c}_2 = \bar{c}_3 = 0$.

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$(y_1, y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = (y_1, y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$0 = -(y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$\begin{aligned} 0 &= && -(y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ z &= && (0 \ 0 \ 2 \ 4)x \end{aligned}$$

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$\begin{aligned} 0 &= && -(y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ z &= && (0 \ 0 \ 2 \ 4)x \end{aligned}$$

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$\begin{aligned} 0 &= && -(y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ z &= && (0 \ 0 \ 2 \ 4)x \end{aligned}$$

$$z = \left[(0 \ 0 \ 2 \ 4) - (y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right] x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$\begin{aligned} 0 &= && -(y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ z &= && (0 \ 0 \ 2 \ 4)x \end{aligned}$$

$$z = \left[(0 \ 0 \ 2 \ 4) - (y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right] x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Remark

For any choice of y_1, y_2 and any feasible solution x ,

$$\max z = (0 \ 0 \ 2 \ 4)x$$

s.t.

$$\begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$x_1, x_2, x_3, x_4 \geq 0$$

$$\begin{aligned} 0 &= && -(y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\ z &= && (0 \ 0 \ 2 \ 4)x \end{aligned}$$

$$z = \left[(0 \ 0 \ 2 \ 4) - (y_1 \ y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right] x + (y_1 \ y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Remark

For any choice of y_1, y_2 and any feasible solution x ,

objective value of x for **old** objective function =
objective value of x for **new** objective function

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^T} x + \underbrace{(y_1 \quad y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}}_{\bar{z}}$$

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^T} x + \underbrace{(y_1 \quad y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}}_{\bar{z}}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_2 = \bar{c}_3 = 0$?

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^\top} x + \underbrace{(y_1 \quad y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}}_{\bar{z}}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_2 = \bar{c}_3 = 0$?

$$(0 \quad 0) = \bar{c}_B^\top = (0 \quad 2) - (y_1 \quad y_2) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^\top} x + \underbrace{(y_1 \quad y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}}_{\bar{z}}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_2 = \bar{c}_3 = 0$?

$$(0 \quad 0) = \bar{c}_B^\top = (0 \quad 2) - (y_1 \quad y_2) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$



$$(y_1 \quad y_2) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} = (0 \quad 2)$$

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^\top} x + \underbrace{(y_1 \quad y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}}_{\bar{z}}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_2 = \bar{c}_3 = 0$?

$$(0 \quad 0) = \bar{c}_B^\top = (0 \quad 2) - (y_1 \quad y_2) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$



$$(y_1 \quad y_2) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} = (0 \quad 2)$$



$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^\top \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \end{pmatrix}$$

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^\top} x + \underbrace{(y_1 \quad y_2) \begin{pmatrix} 1 \\ 2 \end{pmatrix}}_{\bar{z}}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_2 = \bar{c}_3 = 0$?

$$(0 \quad 0) = \bar{c}_B^\top = (0 \quad 2) - (y_1 \quad y_2) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$



$$(y_1 \quad y_2) \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} = (0 \quad 2)$$



$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^\top \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \end{pmatrix}$$



$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 0 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^T} x + \underbrace{(y_1 \quad y_2)}_{\bar{z}} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_B = 0$ for $B = \{2, 3\}$?

Choose

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^T} x + \underbrace{(y_1 \quad y_2)}_{\bar{z}} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_B = 0$ for $B = \{2, 3\}$?

Choose

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$



$$z = \left[(0 \quad 0 \quad 2 \quad 4) - (2 \quad 0) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right] x + (2 \quad 0) \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$z = \underbrace{\left[(0 \quad 0 \quad 2 \quad 4) - (y_1 \quad y_2) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right]}_{\bar{c}^T} x + \underbrace{(y_1 \quad y_2)}_{\bar{z}} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Question

How do we choose y_1, y_2 such that $\bar{c}_B = 0$ for $B = \{2, 3\}$?

Choose

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$



$$z = \left[(0 \quad 0 \quad 2 \quad 4) - (2 \quad 0) \begin{pmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 1 & 2 \end{pmatrix} \right] x + (2 \quad 0) \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$



$$z = (-2 \quad 0 \quad 0 \quad 6)x + 2$$

Rewriting the Objective Function – General

$$\max \quad z = c^\top x$$

s.t.

$$Ax = b$$

$$x \geq 0$$

Rewriting the Objective Function – General

$$\max \quad z = c^\top x$$

s.t.

$$Ax = b$$

$$x \geq 0$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ with $\bar{c}_B = 0$ (\bar{z} constant) for some basis B .

Rewriting the Objective Function – General

$$\max \quad z = c^\top x$$

s.t.

$$Ax = b$$

$$x \geq 0$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ with $\bar{c}_B = 0$ (\bar{z} constant) for some basis B .

$$y^\top Ax = y^\top b$$

Rewriting the Objective Function – General

$$\max \quad z = c^\top x$$

s.t.

$$Ax = b$$

$$x \geq 0$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ with $\bar{c}_B = 0$ (\bar{z} constant) for some basis B .

$$0 = -y^\top Ax + y^\top b$$

Rewriting the Objective Function – General

$$\max \quad z = c^\top x$$

s.t.

$$Ax = b$$

$$x \geq 0$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ with $\bar{c}_B = 0$ (\bar{z} constant) for some basis B .

$$0 = -y^\top Ax + y^\top b$$

$$z = c^\top x$$

Rewriting the Objective Function – General

$$\max \quad z = c^\top x$$

s.t.

$$Ax = b$$

$$x \geq 0$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ with $\bar{c}_B = 0$ (\bar{z} constant) for some basis B .

$$0 = -y^\top Ax + y^\top b$$

$$z = c^\top x$$

$$z = [c^\top - y^\top A] x + y^\top b$$

Rewriting the Objective Function – General

$$\begin{array}{ll} \max & z = c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ with $\bar{c}_B = 0$ (\bar{z} constant) for some basis B .

$$0 = -y^\top Ax + y^\top b$$

$$z = c^\top x$$

$$z = [c^\top - y^\top A] x + y^\top b$$

Remark

For any choice of y and any feasible solution x ,

Rewriting the Objective Function – General

$$\begin{array}{ll} \max & z = c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array}$$

(P2) Replace $c^\top x$ by $\bar{c}^\top x + \bar{z}$ with $\bar{c}_B = 0$ (\bar{z} constant) for some basis B .

$$0 = -y^\top Ax + y^\top b$$

$$z = c^\top x$$

$$z = [c^\top - y^\top A] x + y^\top b$$

Remark

For any choice of y and any feasible solution x ,

$$\begin{array}{l} \text{objective value of } x \text{ for old objective function} \\ \text{objective value of } x \text{ for new objective function} \end{array} =$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$0^\top = \bar{c}_B^\top$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$0^\top = \bar{c}_B^\top = c_B^\top - y^\top A_B$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$0^\top = \bar{c}_B^\top = c_B^\top - y^\top A_B$$



$$y^\top A_B = c_B^\top$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$0^\top = \bar{c}_B^\top = c_B^\top - y^\top A_B$$



$$y^\top A_B = c_B^\top$$



$$A_B^\top y = c_B$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$0^\top = \bar{c}_B^\top = c_B^\top - y^\top A_B$$



$$y^\top A_B = c_B^\top$$



$$A_B^\top y = c_B$$



$$y = (A_B^\top)^{-1} c_B$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$0^\top = \bar{c}_B^\top = c_B^\top - y^\top A_B$$



$$y^\top A_B = c_B^\top$$



$$A_B^\top y = c_B$$



$$y = (A_B^\top)^{-1} c_B$$

Remark

For any non-singular matrix M ,

$$(M^\top)^{-1} = (M^{-1})^\top =: M^{-\top}$$

$$z = \underbrace{[c^\top - y^\top A]}_{\bar{c}^\top} x + \underbrace{y^\top b}_{\bar{z}}$$

Question

How do we choose y such that $\bar{c}_B = 0$ for a basis B ?

$$0^\top = \bar{c}_B^\top = c_B^\top - y^\top A_B$$



$$y^\top A_B = c_B^\top$$



$$A_B^\top y = c_B$$



$$y = (A_B^\top)^{-1} c_B = A_B^{-\top} c_B$$

Remark

For any non-singular matrix M ,

$$(M^\top)^{-1} = (M^{-1})^\top =: M^{-\top}$$

Recap

Proposition

Let B be a basis of A ,

$$\begin{array}{ll} \max & c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array} \quad (\text{P})$$

Recap

Proposition

Let B be a basis of A ,

$$\begin{array}{ll} \max & c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array} \quad (\text{P})$$

$$\begin{array}{ll} \max & \underbrace{[c^\top - y^\top A]}_{\bar{c}} x + y^\top b \\ \text{s.t.} & \\ & \underbrace{A_B^{-1} A}_{A'} x = A_B^{-1} b \\ & x \geq 0 \end{array} \quad (\text{P}')$$

Recap

Proposition

Let B be a basis of A ,

$$\begin{array}{ll} \max & c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array} \quad (\text{P})$$

where $y = A_B^{-\top} c_B$.

$$\begin{array}{ll} \max & \underbrace{[c^\top - y^\top A]}_{\bar{c}} x + y^\top b \\ \text{s.t.} & \\ & \underbrace{A_B^{-1} A}_{A'} x = A_B^{-1} b \\ & x \geq 0 \end{array} \quad (\text{P}')$$

Recap

Proposition

Let B be a basis of A ,

$$\begin{array}{ll} \max & c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array} \quad (\text{P})$$

$$\begin{array}{ll} \max & \underbrace{[c^\top - y^\top A]}_{\bar{c}} x + y^\top b \\ \text{s.t.} & \\ & \underbrace{A_B^{-1} A}_{A'} x = A_B^{-1} b \\ & x \geq 0 \end{array} \quad (\text{P}')$$

where $y = A_B^{-\top} c_B$. Then

(1) (P') is in canonical form for basis B , i.e., $\bar{c}_B = 0$ and $A'_B = I$.

Recap

Proposition

Let B be a basis of A ,

$$\begin{array}{ll} \max & c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array} \quad (\text{P})$$

$$\begin{array}{ll} \max & \underbrace{[c^\top - y^\top A]}_{\bar{c}} x + y^\top b \\ \text{s.t.} & \\ & \underbrace{A_B^{-1} A}_{A'} x = A_B^{-1} b \\ & x \geq 0 \end{array} \quad (\text{P}')$$

where $y = A_B^{-\top} c_B$. Then

- (1) (P') is in canonical form for basis B , i.e., $\bar{c}_B = 0$ and $A'_B = I$.
- (2) (P) and (P') have the same feasible region.

Recap

Proposition

Let B be a basis of A ,

$$\begin{array}{ll} \max & c^\top x \\ \text{s.t.} & \\ & Ax = b \\ & x \geq 0 \end{array} \quad (\text{P})$$

$$\begin{array}{ll} \max & \underbrace{[c^\top - y^\top A]}_{\bar{c}} x + y^\top b \\ \text{s.t.} & \\ & \underbrace{A_B^{-1} A}_{A'} x = A_B^{-1} b \\ & x \geq 0 \end{array} \quad (\text{P}')$$

where $y = A_B^{-\top} c_B$. Then

- (1) (P') is in canonical form for basis B , i.e., $\bar{c}_B = 0$ and $A'_B = I$.
- (2) (P) and (P') have the same feasible region.
- (3) Feasible solutions have the same objective value for (P) and (P').