


Module 2: Linear Programs (Extreme Points)


Extreme Points

Consider the following convex set:




Extreme Points

Consider the following convex set:




Extreme Points

Consider the following convex set:



Extreme Points

Consider the following convex set:



Question

How might we formally describe the “extreme points”?

Towards a Definition of Extreme Points

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if

- $x \in L$ and

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if


- $x \in L$ and
- x is distinct from the endpoints of L .

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if

- $x \in L$ and
- x is distinct from the endpoints of L .



\bar{x} is not contained in L .

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if

- $x \in L$ and
- x is distinct from the endpoints of L .




\bar{x} is contained in L ,
but NOT properly.

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if

- $x \in L$ and
- x is distinct from the endpoints of L .




\bar{x} is PROPERLY contained in L .

Towards a Definition of Extreme Points

Definition

Point $x \in \mathfrak{R}^n$ is **properly contained** in the line segment L if

- $x \in L$ and
- x is distinct from the endpoints of L .



\bar{x} is PROPERLY contained in L .

Definition


Let S be a convex set and $\bar{x} \in S$.

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if

- $x \in L$ and
- x is distinct from the endpoints of L .



\bar{x} is PROPERLY contained in L .

Definition


Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if

Towards a Definition of Extreme Points

Definition

Point $x \in \mathbb{R}^n$ is **properly contained** in the line segment L if

- $x \in L$ and
- x is distinct from the endpoints of L .



\bar{x} is PROPERLY contained in L .

Definition

Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .

Extreme Points - Examples


Definition

Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .

Extreme Points - Examples

Definition


Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .



Extreme Points - Examples

Definition

Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .




Not an extreme point

Extreme Points - Examples

Definition

Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .




Not an extreme point

Extreme Points - Examples

Definition


Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .



Extreme Points - Examples

Definition

Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .




Not an extreme point

Extreme Points - Examples

Definition

Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .




Not an extreme point

Extreme Points - Examples

Definition


Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .



Extreme Points - Examples

Definition

Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .




An extreme point

Extreme Points - Examples

Definition


Let S be a convex set and $\bar{x} \in S$. Then \bar{x} is NOT an **extreme point** if there exists a line segment $L \subseteq S$ where L properly contains \bar{x} .



An extreme point


Question

What are the extreme points in the following figure?




Question

What are the extreme points in the following figure?




Question

What are the extreme points in the following figure?




Question

What are the extreme points in the following figure?




Question

What are the extreme points in the following figure?




Question

What are the extreme points in the following figure?



Question

What are the extreme points in the following figure?




Remark

A convex set may have an **infinite** number of extreme points.


Question

What are the extreme points in the following figure?




Question

What are the extreme points in the following figure?



Question

What are the extreme points in the following figure?



Remark

A convex set may have **NO** extreme points.

This Lecture

This Lecture

Goals:

This Lecture

Goals:

1. Characterize the extreme points in a polyhedron.

This Lecture


Goals:

1. Characterize the extreme points in a polyhedron.
2. Characterize an extreme point for LP in Standard Equality Form.


This Lecture

Goals:

1. Characterize the extreme points in a polyhedron.
2. Characterize an extreme point for LP in Standard Equality Form.
3. Gain a geometric understanding of the Simplex algorithm.



$$P = \left\{ x : \begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} x \leq \begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$




$$P = \left\{ x : \begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} x \leq \begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$

Question

What do the extreme points

$$f = (1, 2)^{\top} \quad \text{and} \quad g = (2, 1)^{\top}$$

have in common?



$$P = \left\{ x : \begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} x \leq \begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$

Question

What do the extreme points

$$f = (1, 2)^\top \quad \text{and} \quad g = (2, 1)^\top$$

have in common?

Each satisfy $n = 2$ “independent” constraints with equality!

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and

Definition


Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.




$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right\} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix}$$

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.




Consider f :

$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right\} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix}$$

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.



Consider f :


Here (1) and (3) are tight.

$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right\} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix}$$

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right\} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix}$$


Consider f :

Here (1) and (3) are tight. $\underbrace{\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}}_{\bar{A}} x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \end{pmatrix}}_{\bar{b}}$.

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.




Consider g :

$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right\} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix}$$

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.



Consider g :

Here (1) and (2) are tight.


$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right\}$$

(1)
(2)
(3)

Definition

Let $P = \{x : Ax \leq b\}$ be a polyhedron and let $x \in P$.

- A constraint is **tight** for x if it is satisfied with equality, and
- the set of all tight constraints is denoted $\bar{A}x \leq \bar{b}$.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right\} \begin{matrix} (1) \\ (2) \\ (3) \end{matrix}$$

Consider g :

Here (1) and (2) are tight. $\underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}}_{\bar{A}} x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \end{pmatrix}}_{\bar{b}}$.

Theorem


Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.




$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.




Consider f :

$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right. \left. \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$


Consider f :

$$\bar{A} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right. \left. \begin{array}{l} (1) \\ (2) \\ (3) \end{array} \right\}$$


Consider f :

$\bar{A} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ so, since $\text{rank}(\bar{A}) = 2$, f is an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.




$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.




Consider g :

$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right. \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \left. \right\}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right. \left. \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$


Consider g :

$$\bar{A} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right. \left. \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$


Consider g :

$\bar{A} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ so, since $\text{rank}(\bar{A}) = 2$, g is an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.




$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.




Consider h :

$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \begin{matrix} (1) \\ (2) \\ (3) \end{matrix} \right\}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right. \left. \begin{array}{l} (1) \\ (2) \\ (3) \end{array} \right\}$$


Consider h :

$$\bar{A} = (1 \ 0)$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}}_b \right. \left. \begin{array}{l} (1) \\ (2) \\ (3) \end{array} \right\}$$

Consider h :

$\bar{A} = \begin{pmatrix} 1 & 0 \end{pmatrix}$ so, since $\text{rank}(\bar{A}) < 2$, h is NOT an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Is the following true?

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Is the following true?

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Is the following true? **NO!**

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Is the following true? **NO!**


Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Is the following true? **NO!**

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.




$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 0 \\ 0 \end{pmatrix}}_b \right\}$$

Is the following true? **NO!**

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.



$$P = \left\{ x : \underbrace{\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}}_A x \leq \underbrace{\begin{pmatrix} 0 \\ 0 \end{pmatrix}}_b \right\}$$

$\bar{A} = \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}$ has $n = 2$ rows, but $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ is NOT extreme.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Let's prove part (1).

Remark

Let $a, b, c \in \mathbb{R}$, and

Remark

Let $a, b, c \in \mathbb{R}$, and suppose

$$a = \frac{1}{2}b + \frac{1}{2}c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Remark

Let $a, b, c \in \mathbb{R}$, and suppose

$$a = \frac{1}{2}b + \frac{1}{2}c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Then $a = b = c$.

Remark

Let $a, b, c \in \mathbb{R}$, and suppose

$$a = \frac{1}{2}b + \frac{1}{2}c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Then $a = b = c$.

Proof

$$a = \frac{1}{2} b + \frac{1}{2} c$$

Remark

Let $a, b, c \in \mathbb{R}$, and suppose

$$a = \frac{1}{2}b + \frac{1}{2}c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Then $a = b = c$.

Proof

$$a = \frac{1}{2} \underbrace{b}_{\leq a} + \frac{1}{2} \underbrace{c}_{\leq a}$$

Remark

Let $a, b, c \in \mathbb{R}$, and suppose

$$a = \frac{1}{2}b + \frac{1}{2}c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Then $a = b = c$.

Proof

$$a = \frac{1}{2} \underbrace{b}_{\leq a} + \frac{1}{2} \underbrace{c}_{\leq a} \leq \frac{1}{2}a + \frac{1}{2}a = a.$$

Remark

Let $a, b, c \in \mathbb{R}$, and suppose

$$a = \frac{1}{2}b + \frac{1}{2}c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Then $a = b = c$.

Proof

$$a = \frac{1}{2} \underbrace{b}_{\leq a} + \frac{1}{2} \underbrace{c}_{\leq a} \leq \frac{1}{2}a + \frac{1}{2}a = a.$$

Thus, equality holds throughout

Remark

Let $a, b, c \in \mathbb{R}$, and suppose

$$a = \frac{1}{2}b + \frac{1}{2}c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Then $a = b = c$.

Proof

$$a = \frac{1}{2} \underbrace{b}_{\leq a} + \frac{1}{2} \underbrace{c}_{\leq a} \leq \frac{1}{2}a + \frac{1}{2}a = a.$$

Thus, equality holds throughout $\Rightarrow b = a$ and $c = a$.

Remark

Let $a, b, c \in \mathbb{R}$,

Remark

Let $a, b, c \in \mathbb{R}$, and let λ where $0 < \lambda < 1$.

Remark

Let $a, b, c \in \mathbb{R}$, and let λ where $0 < \lambda < 1$. Suppose

$$a = \lambda b + (1 - \lambda)c \quad \text{and} \quad b \leq a, c \leq a.$$

Remark

Let $a, b, c \in \mathbb{R}$, and let λ where $0 < \lambda < 1$. Suppose

$$a = \lambda b + (1 - \lambda)c \quad \text{and} \quad b \leq a, \quad c \leq a.$$

Then $a = b = c$.

Remark

Let $a, b, c \in \mathbb{R}$, and let λ where $0 < \lambda < 1$. Suppose

$$a = \lambda b + (1 - \lambda)c \quad \text{and} \quad b \leq a, c \leq a.$$

Then $a = b = c$.

Exercise

Prove the previous remark.

Remark

Let $a, b, c \in \mathbb{R}^n$

Remark

Let $a, b, c \in \mathbb{R}^n$, and let λ where $0 < \lambda < 1$. Suppose

Remark

Let $a, b, c \in \mathbb{R}^n$, and let λ where $0 < \lambda < 1$. Suppose

$$a = \lambda b + (1 - \lambda)c \quad \text{and} \quad b \leq a, c \leq a.$$

Remark

Let $a, b, c \in \mathbb{R}^n$, and let λ where $0 < \lambda < 1$. Suppose

$$a = \lambda b + (1 - \lambda)c \quad \text{and} \quad b \leq a, c \leq a.$$

Then $a = b = c$.

Remark

Let $a, b, c \in \mathbb{R}^n$, and let λ where $0 < \lambda < 1$. Suppose

$$a = \lambda b + (1 - \lambda)c \quad \text{and} \quad b \leq a, c \leq a.$$

Then $a = b = c$.

Exercise

Prove the previous remark.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$,

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

$$\bar{b} = \bar{A}\bar{x}$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

$$\bar{b} = \bar{A}\bar{x} = \bar{A} \left(\lambda x^{(1)} + (1 - \lambda)x^{(2)} \right)$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

$$\bar{b} = \bar{A}\bar{x} = \bar{A} \left(\lambda x^{(1)} + (1 - \lambda)x^{(2)} \right) = \lambda \bar{A}x^{(1)} + (1 - \lambda)\bar{A}x^{(2)}.$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

$$\bar{b} = \bar{A}\bar{x} = \bar{A}(\lambda x^{(1)} + (1 - \lambda)x^{(2)}) = \lambda \bar{A}x^{(1)} + (1 - \lambda)\bar{A}x^{(2)}.$$

$\bar{A}x^{(1)} \leq \bar{b}$ and $\bar{A}x^{(2)} \leq \bar{b}$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

$$\bar{b} = \bar{A}\bar{x} = \bar{A}(\lambda x^{(1)} + (1 - \lambda)x^{(2)}) = \lambda \bar{A}x^{(1)} + (1 - \lambda)\bar{A}x^{(2)}.$$

$\bar{A}x^{(1)} \leq \bar{b}$ and $\bar{A}x^{(2)} \leq \bar{b}$.

Previous remark implies that $\bar{b} = \bar{A}x^{(1)} = \bar{A}x^{(2)}$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

$$\bar{b} = \bar{A}\bar{x} = \bar{A} \left(\lambda x^{(1)} + (1 - \lambda)x^{(2)} \right) = \lambda \bar{A}x^{(1)} + (1 - \lambda)\bar{A}x^{(2)}.$$

$$\bar{A}x^{(1)} \leq \bar{b} \text{ and } \bar{A}x^{(2)} \leq \bar{b}.$$

Previous remark implies that $\bar{b} = \bar{A}x^{(1)} = \bar{A}x^{(2)}$.

However, since $\text{rank}(\bar{A}) = n$, $x^{(1)} = x^{(2)}$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.

Proof

Suppose \bar{x} is not an extreme point.

\bar{x} is properly contained in a line segment with endpoints $x^{(1)}, x^{(2)} \in P$.

$\bar{x} \neq x^{(1)}, x^{(2)} \in P$ and for some λ , $0 < \lambda < 1$, $\bar{x} = \lambda x^{(1)} + (1 - \lambda)x^{(2)}$.

$$\bar{b} = \bar{A}\bar{x} = \bar{A}(\lambda x^{(1)} + (1 - \lambda)x^{(2)}) = \lambda \bar{A}x^{(1)} + (1 - \lambda)\bar{A}x^{(2)}.$$

$$\bar{A}x^{(1)} \leq \bar{b} \text{ and } \bar{A}x^{(2)} \leq \bar{b}.$$

Previous remark implies that $\bar{b} = \bar{A}x^{(1)} = \bar{A}x^{(2)}$.

However, since $\text{rank}(\bar{A}) = n$, $x^{(1)} = x^{(2)}$. This is a contradiction.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Proof

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$.

$$x^{(1)} = \bar{x} + \epsilon d$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$.

$$x^{(1)} = \bar{x} + \epsilon d$$

$$x^{(2)} = \bar{x} - \epsilon d$$

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.


Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$.

$$x^{(1)} = \bar{x} + \epsilon d$$

$$x^{(2)} = \bar{x} - \epsilon d$$



Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Proof


Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = 0$.

Pick a **small** $\epsilon > 0$.

$$x^{(1)} = \bar{x} + \epsilon d$$

$$x^{(2)} = \bar{x} - \epsilon d$$

It suffices to prove the following:



Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.


Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$.

$$x^{(1)} = \bar{x} + \epsilon d$$

$$x^{(2)} = \bar{x} - \epsilon d$$



It suffices to prove the following:

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.


Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$.

$$x^{(1)} = \bar{x} + \epsilon d$$

$$x^{(2)} = \bar{x} - \epsilon d$$



It suffices to prove the following:

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

(a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.

Why?

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

(a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.

Why?

$$\frac{1}{2}x^{(1)} + \frac{1}{2}x^{(2)}$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

(a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.

Why?

$$\frac{1}{2}x^{(1)} + \frac{1}{2}x^{(2)} = \frac{1}{2}(\bar{x} + \epsilon d) + \frac{1}{2}(\bar{x} - \epsilon d)$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

(a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.

Why?

$$\frac{1}{2}x^{(1)} + \frac{1}{2}x^{(2)} = \frac{1}{2}(\bar{x} + \epsilon d) + \frac{1}{2}(\bar{x} - \epsilon d) = \bar{x}.$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

(a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.

Why?

$$\frac{1}{2}x^{(1)} + \frac{1}{2}x^{(2)} = \frac{1}{2}(\bar{x} + \epsilon d) + \frac{1}{2}(\bar{x} - \epsilon d) = \bar{x}. \quad \checkmark$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

(a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)}$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d)$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}}$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}} = \bar{b}.$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}} = \bar{b}. \quad \checkmark$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}} = \bar{b}. \quad \checkmark$$

Consider non-tight constraint $a^\top x \leq \beta$.

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}} = \bar{b}. \quad \checkmark$$

Consider non-tight constraint $a^\top x \leq \beta$.

$$a^\top x^{(1)}$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}} = \bar{b}. \quad \checkmark$$

Consider non-tight constraint $a^\top x \leq \beta$.

$$a^\top x^{(1)} = a^\top (\bar{x} + \epsilon d)$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}} = \bar{b}. \quad \checkmark$$

Consider non-tight constraint $a^\top x \leq \beta$.

$$a^\top x^{(1)} = a^\top (\bar{x} + \epsilon d) = \underbrace{a^\top \bar{x}}_{< \beta} + \epsilon \underbrace{a^\top d}_{??}$$

Proof

Since $\text{rank}(\bar{A}) < n$, there exists a **non-zero** vector d such that $\bar{A}d = \mathbf{0}$.

Pick a **small** $\epsilon > 0$. Let $x^{(1)} = \bar{x} + \epsilon d$ and $x^{(2)} = \bar{x} - \epsilon d$.

- (a) \bar{x} is properly contained in the line segment between $x^{(1)}$ and $x^{(2)}$.
- (b) $x^{(1)}, x^{(2)} \in P$. (It is sufficient to show this for $x^{(1)}$ only.)

Consider tight constraints $\bar{A}x \leq \bar{b}$.

$$\bar{A}x^{(1)} = \bar{A}(\bar{x} + \epsilon d) = \underbrace{\bar{A}\bar{x}}_{\bar{b}} + \epsilon \underbrace{\bar{A}d}_{\mathbf{0}} = \bar{b}. \quad \checkmark$$

Consider non-tight constraint $a^\top x \leq \beta$.

$$a^\top x^{(1)} = a^\top (\bar{x} + \epsilon d) = \underbrace{a^\top \bar{x}}_{< \beta} + \epsilon \underbrace{a^\top d}_{??} < \beta$$

for a **small enough** ϵ .

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\}$$

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

Let's use our theorem to find an answer.

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

Let's use our theorem to find an answer.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

Let's use our theorem to find an answer.

Theorem

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron and let $\bar{x} \in P$.

1. If $\text{rank}(\bar{A}) = n$, then \bar{x} is an extreme point.
2. If $\text{rank}(\bar{A}) < n$, then \bar{x} is NOT an extreme point.

We need to rewrite the constraints in P so they are all in the form " \leq ".

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

We need to rewrite the constraints in P so they are all in the form " \leq ".

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

We need to rewrite the constraints in P so they are all in the form " \leq ".

$P = \{x : Ax \leq b\}$, where

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline & & \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 2 \\ 4 \\ \hline \end{pmatrix}$$

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

We need to rewrite the constraints in P so they are all in the form " \leq ".

$P = \{x : Ax \leq b\}$, where

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 2 \\ 4 \\ \hline -2 \\ -4 \\ \hline \end{pmatrix}$$

Consider

$$P = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \end{pmatrix} x = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\} \quad \begin{pmatrix} 2 \\ 4 \\ 0 \end{pmatrix} \text{ is a basic solution}$$

Question

Is $(2, 4, 0)^\top$ an extreme point?

We need to rewrite the constraints in P so they are all in the form " \leq ".

$P = \{x : Ax \leq b\}$, where

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 2 \\ 4 \\ \hline -2 \\ -4 \\ \hline 0 \\ 0 \\ 0 \end{pmatrix}$$

For $P = \{x : Ax \leq b\}$, where

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad b = \begin{pmatrix} 2 \\ 4 \\ \hline -2 \\ -4 \\ \hline 0 \\ 0 \\ 0 \end{pmatrix}$$

and $(2, 4, 0)^\top$, we have

For $P = \{x : Ax \leq b\}$, where

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad b = \begin{pmatrix} 2 \\ 4 \\ \hline -2 \\ -4 \\ \hline 0 \\ 0 \\ 0 \end{pmatrix}$$

and $(2, 4, 0)^\top$, we have

$$\bar{A} = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline 0 & 0 & -1 \end{pmatrix}$$

For $P = \{x : Ax \leq b\}$, where

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad b = \begin{pmatrix} 2 \\ 4 \\ \hline -2 \\ -4 \\ \hline 0 \\ 0 \\ 0 \end{pmatrix}$$

and $(2, 4, 0)^\top$, we have

$$\bar{A} = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline 0 & 0 & -1 \end{pmatrix}$$

Since $\text{rank}(\bar{A}) = 3$, we know that $(2, 4, 0)^\top$ is an extreme point!

For $P = \{x : Ax \leq b\}$, where

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad b = \begin{pmatrix} 2 \\ 4 \\ \hline -2 \\ -4 \\ \hline 0 \\ 0 \\ 0 \end{pmatrix}$$

and $(2, 4, 0)^\top$, we have

$$\bar{A} = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 3 \\ \hline -1 & 0 & 1 \\ 0 & -1 & -3 \\ \hline 0 & 0 & -1 \end{pmatrix}$$

Since $\text{rank}(\bar{A}) = 3$, we know that $(2, 4, 0)^\top$ is an extreme point!

This is no accident...

Theorem

Let $P = \{x \geq \mathbf{0} : Ax = b\}$ where rows of A are independent. The following are equivalent:

1. \bar{x} is an extreme point of P .
2. \bar{x} is a basic feasible solution of P .

Theorem

Let $P = \{x \geq \mathbf{0} : Ax = b\}$ where rows of A are independent. The following are equivalent:

1. \bar{x} is an extreme point of P .
2. \bar{x} is a basic feasible solution of P .

Exercise

Prove the previous theorem.

Theorem

Let $P = \{x \geq \mathbf{0} : Ax = b\}$ where rows of A are independent. The following are equivalent:

1. \bar{x} is an extreme point of P .
2. \bar{x} is a basic feasible solution of P .

Exercise

Prove the previous theorem.



Theorem

Let $P = \{x \geq \mathbf{0} : Ax = b\}$ where rows of A are independent. The following are equivalent:

1. \bar{x} is an extreme point of P .
2. \bar{x} is a basic feasible solution of P .

Exercise

Prove the previous theorem.



The Simplex algorithm moves from extreme points to extreme points.

Simplex - a Geometric Illustration

Simplex - a Geometric Illustration

$$\max (2, 3, 0, 0, 0)x$$

s.t.

$$x \in P_1$$

Simplex - a Geometric Illustration

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Simplex - a Geometric Illustration

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$

Simplex - a Geometric Illustration

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$:

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$: **optimal**

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$: **optimal**



Simplex visits **extreme points** of P_1 in order:

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$: **optimal**

 Simplex visits **extreme points** of P_1 in order:

$$\begin{pmatrix} 0 \\ 0 \\ 10 \\ 6 \\ 4 \end{pmatrix},$$

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$: **optimal**

 Simplex visits **extreme points** of P_1 in order:

$$\begin{pmatrix} 0 \\ 0 \\ 10 \\ 6 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \\ 0 \\ 1 \\ 9 \end{pmatrix},$$

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$: **optimal**

 Simplex visits **extreme points** of P_1 in order:

$$\begin{pmatrix} 0 \\ 0 \\ 10 \\ 6 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \\ 0 \\ 1 \\ 9 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \\ 0 \\ 0 \\ 6 \end{pmatrix},$$

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$: **optimal**

 Simplex visits **extreme points** of P_1 in order:

$$\begin{pmatrix} 0 \\ 0 \\ 10 \\ 6 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \\ 0 \\ 1 \\ 9 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \\ 0 \\ 0 \\ 6 \end{pmatrix}, \begin{pmatrix} 1 \\ 5 \\ 3 \\ 0 \\ 0 \end{pmatrix}.$$

Simplex - a Geometric Illustration

$$\begin{array}{l} \max \quad (2, 3, 0, 0, 0)x \\ \text{s.t.} \\ \quad x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

SOLVE USING SIMPLEX:

- Basis $B = \{3, 4, 5\}$, basic solution $(0, 0, 10, 6, 4)^\top$
- Basis $B = \{1, 4, 5\}$, basic solution $(5, 0, 0, 1, 9)^\top$
- Basis $B = \{1, 2, 5\}$, basic solution $(4, 2, 0, 0, 6)^\top$
- Basis $B = \{1, 2, 3\}$, basic solution $(1, 5, 3, 0, 0)^\top$: **optimal**

 Simplex visits **extreme points** of P_1 in order:

$$\begin{pmatrix} 0 \\ 0 \\ 10 \\ 6 \\ 4 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \\ 0 \\ 1 \\ 9 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \\ 0 \\ 0 \\ 6 \end{pmatrix}, \begin{pmatrix} 1 \\ 5 \\ 3 \\ 0 \\ 0 \end{pmatrix}.$$

However, we cannot draw a picture of this...

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Remark

$(0, 0, 10, 6, 4)^\top$ extreme point of $P_1 \Rightarrow (0, 0)^\top$ extreme point of P_2 ,

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Remark

$(0, 0, 10, 6, 4)^\top$	extreme point of P_1	\Rightarrow	$(0, 0)^\top$	extreme point of P_2 ,
$(5, 0, 0, 1, 9)^\top$	extreme point of P_1	\Rightarrow	$(5, 0)^\top$	extreme point of P_2 ,
$(4, 2, 0, 0, 6)^\top$	extreme point of P_1	\Rightarrow	$(4, 2)^\top$	extreme point of P_2 ,
$(1, 5, 3, 0, 0)^\top$	extreme point of P_1	\Rightarrow	$(1, 5)^\top$	extreme point of P_2 .

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Remark

$(0, 0, 10, 6, 4)^\top$	extreme point of P_1	\Rightarrow	$(0, 0)^\top$	extreme point of P_2 ,
$(5, 0, 0, 1, 9)^\top$	extreme point of P_1	\Rightarrow	$(5, 0)^\top$	extreme point of P_2 ,
$(4, 2, 0, 0, 6)^\top$	extreme point of P_1	\Rightarrow	$(4, 2)^\top$	extreme point of P_2 ,
$(1, 5, 3, 0, 0)^\top$	extreme point of P_1	\Rightarrow	$(1, 5)^\top$	extreme point of P_2 .



Simplex visits **extreme points** of P_2 in order:

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Remark

$(0, 0, 10, 6, 4)^\top$	extreme point of P_1	\Rightarrow	$(0, 0)^\top$	extreme point of P_2 ,
$(5, 0, 0, 1, 9)^\top$	extreme point of P_1	\Rightarrow	$(5, 0)^\top$	extreme point of P_2 ,
$(4, 2, 0, 0, 6)^\top$	extreme point of P_1	\Rightarrow	$(4, 2)^\top$	extreme point of P_2 ,
$(1, 5, 3, 0, 0)^\top$	extreme point of P_1	\Rightarrow	$(1, 5)^\top$	extreme point of P_2 .



Simplex visits **extreme points** of P_2 in order:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Remark

$(0, 0, 10, 6, 4)^\top$	extreme point of P_1	\Rightarrow	$(0, 0)^\top$	extreme point of P_2 ,
$(5, 0, 0, 1, 9)^\top$	extreme point of P_1	\Rightarrow	$(5, 0)^\top$	extreme point of P_2 ,
$(4, 2, 0, 0, 6)^\top$	extreme point of P_1	\Rightarrow	$(4, 2)^\top$	extreme point of P_2 ,
$(1, 5, 3, 0, 0)^\top$	extreme point of P_1	\Rightarrow	$(1, 5)^\top$	extreme point of P_2 .



Simplex visits **extreme points** of P_2 in order:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \end{pmatrix},$$

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Remark

$(0, 0, 10, 6, 4)^\top$	extreme point of P_1	\Rightarrow	$(0, 0)^\top$	extreme point of P_2 ,
$(5, 0, 0, 1, 9)^\top$	extreme point of P_1	\Rightarrow	$(5, 0)^\top$	extreme point of P_2 ,
$(4, 2, 0, 0, 6)^\top$	extreme point of P_1	\Rightarrow	$(4, 2)^\top$	extreme point of P_2 ,
$(1, 5, 3, 0, 0)^\top$	extreme point of P_1	\Rightarrow	$(1, 5)^\top$	extreme point of P_2 .



Simplex visits **extreme points** of P_2 in order:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \end{pmatrix},$$

$$\begin{array}{ll} \max & (2, 3, 0, 0, 0)x \\ \text{s.t.} & \\ & x \in P_1 \end{array}$$

$$P_1 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 1 \end{pmatrix} x = \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

is obtained by adding **slack variables** to

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Remark

$(0, 0, 10, 6, 4)^\top$	extreme point of P_1	\Rightarrow	$(0, 0)^\top$	extreme point of P_2 ,
$(5, 0, 0, 1, 9)^\top$	extreme point of P_1	\Rightarrow	$(5, 0)^\top$	extreme point of P_2 ,
$(4, 2, 0, 0, 6)^\top$	extreme point of P_1	\Rightarrow	$(4, 2)^\top$	extreme point of P_2 ,
$(1, 5, 3, 0, 0)^\top$	extreme point of P_1	\Rightarrow	$(1, 5)^\top$	extreme point of P_2 .



Simplex visits **extreme points** of P_2 in order:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 5 \end{pmatrix}.$$

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$


$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Simplex visits extreme points of P_2 in order: $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 5 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$, $\begin{pmatrix} 1 \\ 5 \end{pmatrix}$.

$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$


Simplex visits extreme points of P_2 in order: $\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 5 \end{pmatrix}$.



$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$


Simplex visits extreme points of P_2 in order: $\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 5 \\ 0 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 5 \end{pmatrix}$.



$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$


Simplex visits **extreme points** of P_2 in order: $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\boxed{\begin{pmatrix} 5 \\ 0 \end{pmatrix}}$, $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$, $\begin{pmatrix} 1 \\ 5 \end{pmatrix}$.



$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$


Simplex visits **extreme points** of P_2 in order: $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 5 \\ 0 \end{pmatrix}$, $\boxed{\begin{pmatrix} 4 \\ 2 \end{pmatrix}}$, $\begin{pmatrix} 1 \\ 5 \end{pmatrix}$.



$$\begin{array}{ll} \max & (2, 3)x \\ \text{s.t.} & \\ & x \in P_2 \end{array}$$

$$P_2 = \left\{ x \geq \mathbf{0} : \begin{pmatrix} 2 & 1 \\ 1 & 1 \\ -1 & 1 \end{pmatrix} x \leq \begin{pmatrix} 10 \\ 6 \\ 4 \end{pmatrix} \right\}$$

Simplex visits **extreme points** of P_2 in order: $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 5 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$, $\boxed{\begin{pmatrix} 1 \\ 5 \end{pmatrix}}$.



Recap

Recap

- We defined extreme points of convex sets.

Recap

- We defined extreme points of convex sets.
- We characterized extreme points in polyhedra.

Recap

- We defined extreme points of convex sets.
- We characterized extreme points in polyhedra.
- We saw that extreme points = basic solutions for problems in SEF.

Recap

- We defined extreme points of convex sets.
- We characterized extreme points in polyhedra.
- We saw that extreme points = basic solutions for problems in SEF.
- We showed that the Simplex algorithm moves from extreme point to extreme point.